



# **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
04.09.2002 Bulletin 2002/36

(51) Int Cl.7: **C12N 15/56, C12N 9/24, C12N 15/80, C12Q 1/68, C12P 19/14, C11D 3/386**

(21) Application number: **92902669.8**

(86) International application number:  
**PCT/US91/09285**

(22) Date of filing: **10.12.1991**

(87) International publication number:  
**WO 92/010581 (25.06.1992 Gazette 1992/14)**

(54) **IMPROVED SACCHARIFICATION OF CELLULOSE BY CLONING AND AMPLIFICATION OF THE BETA-GLUCOSIDASE GENE OF TRICHODERMA REESEI**

VERBESSERTE SACCHARIFIZIERUNG VON ZELLULOSE DURCH KLONIERUNG UND VERVIELFÄLTIGUNG DES BETA-GLUKOSIDASE GENES AUS TRICHODERMA REESEI

SACCHARIFICATION AMELIOREE DE CELLULOSE PAR CLONAGE ET AMPLIFICATION DU GENE DE BETA-GLUCOSIDASE DE TRICHODERMA REESEI

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU MC NL SE**

(30) Priority: **10.12.1990 US 625140**

(43) Date of publication of application:  
**29.09.1993 Bulletin 1993/39**

(60) Divisional application:  
**02005972.1 / 1 225 227**

(73) Proprietor: **GENENCOR INTERNATIONAL, INC.**  
**Palo Alto, California 94304 (US)**

(72) Inventors:  
• **FOWLER, Timothy**  
**Belmont, CA 94002 (US)**  
• **BARNETT, Christopher, C.**  
**South San Francisco, CA 94080 (US)**  
• **SHOEMAKER, Sharon**  
**Fairfield, CA 94533 (US)**

(74) Representative: **Armitage, Ian Michael et al**  
**MEWBURN ELLIS**  
**York House**  
**23 Kingsway**  
**London WC2B 6HP (GB)**

(56) References cited:

<b>EP-A- 0 148 668</b>	<b>WO-A-91/17244</b>
<b>WO-A-92/06209</b>	<b>WO-A-92/06210</b>
<b>DD-A- 263 571</b>	<b>US-A- 4 275 163</b>
<b>US-A- 4 472 504</b>	<b>US-A- 4 885 252</b>
<b>US-A- 4 892 819</b>	<b>US-A- 4 935 349</b>

- **DATABASE WPI Derwent Publications Ltd., London, GB; AN 88-305166 & JP-A-63 226 294 (AJINIMOTO KK) 20 September 1988**
- **DATABASE WPI Derwent Publications Ltd., London, GB; AN 93-190701 & JP-A-5 115 293 (FOOD DESIGN GIJUTSU KENKYUKUMIAI) 14 May 1993**
- **Proceedings of the National Science Council Republic of China, Part B life Science, Vol. 12, No. 2, issued 1988, C.H. WANG et al., "An efficient approach for the preparation of fungal protoplasts: protoplast formation and regeneration in Trichoderma kiningii, pages 69-76, see Abstract No. 6960172.**
- **Hoppe-Seyler's Zeischrift fuer Physiologische Chemie, Vol. 355, No. 4, issued 1974, E. BAUSE, "Isolation and amino acid sequence of a hexa deca peptide from the active site of beta-glucosidase A-3 from Aspergillus wentii", pages 438-442, see Abstract No. 1585209.**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 562 003 B1**

- Molecular General Genetics, Vol. 217, issued May 1989, W.L. STAUDENBAUER, "Nucleotide sequence of the *Clostridium thermocellum* bg1B gene encoding thermally stable beta-glucosidase: homology to fungal beta-glucosidase", pages 70-76, see entire document.
- Journal of Applied Biochemistry, Vol. 6, issued 1984, P. BHIKHABHAI et al., "Isolation of cellulolytic enzymes from *Trichoderma reesei*", pages 336-345, see Abstract No. 05644951.
- *Trichoderma reesei* Cellulases, issued 1990, H. MERIVUORI et al., "Effects of alcohol and temperature on secretion by *Trichoderma reesei*: cellulase complex production by *Trichoderma reesei*; influenza A virus hemagglutinin gene cloning and expression", pages 105-114, see Abstract No. 91-09562.
- Bio/Technology, Vol. 9, issued June 1991, C.C. BARNETT, "Cloning and amplification of the gene encoding and extracellular beta-glucosidase from *Trichoderma reesei*: evidence for improved rates of saccharification of cellulosic substrates: DNA sequence", pages 562-566.
- Protein Engineering, Vol. 3, no. 4, issued 1990, R.C. MILLER, Jr. et al., "Enzyme mobilization or purification using the cellulose-binding domains of the two bacterial cellulases - *Cellulomonas fimi* cellobiohydrolase and cellulase cellulose-binding domain, *Aspergillus* sp. beta-glucosidase, construction", page 379, see Abstract No. 90-06574.
- Critical Rev. Biotechnol., Vol. 9, Vol. 2, issued 1989, V.S. BISARIA et al., "Regulatory aspects of cellulase biosynthesis and secretion: beta-glucosidase protein secretion", pages 64-103, see Abstract no. 90-04716.
- Molecular General Genetics, Vol. 194, issued 1984, M.E. Penttila et al., "Cloning of *Aspergillus niger* genes in yeast. Expression of the gene coding *Aspergillus* beta-glucosidase", pages 494-499, see entire document.
- Current Genetics, Vol. 12, issued 1987, A. RAYNAL et al., "Sequence and transcription of the beta-glucosidase gene of *Kluyveromyces fragilis* in *Saccharomyces cerevisiae*", pages 175-184, see entire document.
- Journal of Applied Bacteriology, Vol. 61, issued 1986, M.K. KALRA et al., "Partial purification, characterization and regulation of cellulolytic enzymes from *Trichoderma longibrachiatum*", pages 73-80, see entire document.
- Proceedings of the National Academy of Sciences U.S.A. Vol. 82, issued 1985, M.M. YELTON et al., "A cosmid for selecting genes by complementation in *Aspergillus nidulans*: selection of the developmentally regulated *yA* locus", pages 834-838, see entire document.
- Biochimica et Biophysica Acta, Vol. 626, issued 1980, ERNST BAUSE et al., "Isolation and structure of a tryptic glycopeptide from the active site of beta-glucosidase A3 from *Aspergillus wentii*", pages 459-465.
- Journal of Bacteriology, Vol. 170, No. 1, issued January 1988, W.W. WAKARCHUK et al., "Structure and transcription analysis of the gene encoding a cellobiose from *Agrobacterium* sp. strain ATCC 21400", pages 301-307, see entire document.
- Current Genetics, Vol. 12, issued 1987, M.E. PENTTILA et al., "Construction of brewer's yeast secreting fungal endo-beta-glucanase", pages 413-420, see entire document.
- FEMS Microbiology Letters, Vol. 30, issued 1985, N.A. SAHASRABUDHE et al., "Genome complexity of a powerful cellulolytic fungus, *Penicillium funiculosum*", pages 295-300.
- FEBS Congress 16th Meeting, Part C, issued 1985, J. KNOWLES et al., "The cloning of fungal cellulase genes and their expression in yeast: cellobiohydrolase and beta-glucosidase gene expression in transformed *Saccharomyces cerevisiae*", pages 43-49, see Abstract No. 86-06126.
- Biotechnol. Adv., Vol. 7, No. 3, issued 1989, B.R. GLICK et al., "Isolation, characterization and manipulation of cellulase genes: cellobiohydrolase and beta-glucosidase genes cloning and expression; cellulase complex; a review", pages 361-386, see Abstract No. 90-02276.
- Abstr. Pap. Am. Chem. Soc., 194 Meeting, issued 1987, M. GRITZALI, "Enzymes of the cellulase system of *Trichoderma*: an overview - cellulase complex characterization", see Abstract No. 88-04459.
- FEMS Symp., Vol. 43, issued 1988, H. DURAND et al., "Classical and molecular genetics applied to *Trichoderma reesei* from the selection of improved cellulolytic industrial strains - beta-glucosidase and cellulase activity; beta-D-fructofuranosidase gene cloning and expression in *Trichoderma reesei*; cellulose degradation", see Abstract No. 89-11621.
- Abstr. Pap. Am. Chem. Soc., 195 Meeting, issued 1988, C. BARNETT et al., "Expression of *Trichoderma reesei* exo-cellobiohydrolase II genes in *Aspergillus awamori*: a heterologous expression system to study structure-function relationships - enzyme engineering applications", see Abstract No. 88-08174.
- FEMS Symp., Vol. 43, issued 1988, J. KNOWLES et al., "The use of gene technology to investigate fungal cellulolytic enzymes - *Trichoderma reesei* cellulase complex gene cloning and expression in *Saccharomyces cerevisiae*", see Abstract No. 89-11622.

- World Biotech. Rep., Vol. 2, issued 1984, S.P. SHOEMAKER et al., "The cellulase system of *Trichoderma reesei*: *Trichoderma* strain improvement and expression of *Trichoderma cellulase* in yeast -- in ethanol production", pages 593-600, see Abstract No. 85-02556.
- Biochem. Soc. Trans., Vol. 13, No. 2, issued 1985, K. MERIVUORI et al., "Regulation of cellulase biosynthesis and secretion in fungi -- *Trichoderma reesei*", pages 411-414, see Abstract No. 85-11852.
- Abstr. Pap. Am. Chem. Soc., 202 Meeting, issued 1991, T. FOWLER, "Developments in recombinant beta-glucosidase of *Trichoderma reesei* -- cellulase complex recombinant beta-glucosidase production", see Abstract No. 91-14368.
- Biotechnol., Vol. 7, No. 9, issued 1989, ONG et al., "The cellulose-binding domains of cellulases: to facilitate biotechnology-application in protein or fusel protein purification by affinity chromatography and in enzyme immobilization, a review", pages 239-243, see Abstract No. 89-12873.
- Aust. J. Biotechnol., Vol. 3, No. 1, issued 1989, P.A.D. RICKARD et al., "Kinetic properties and contribution to cellulose saccharification of a cloned *Pseudomonas* beta-glucosidase: enzyme cloning using plasmid pND71 and use of recombinant product in association with *Trichoderma reesei* cellulase", pages 43-49, see Abstract No. 90-11472.

## Description

## BACKGROUND OF THE INVENTION

## 5 1. Field of the Invention.

[0001] The present invention relates to cellulase preparations and compositions having increased or decreased cellulolytic capacity. The invention further relates to a nucleotide sequence of the bgl1 gene encoding extracellular  $\beta$ -glucosidase from a filamentous fungi, a plasmid vector containing the gene encoding extracellular  $\beta$ -glucosidase and transformant strains with increased copy numbers of the  $\beta$ -glucosidase (bgl1) gene introduced into the genome. More particularly, the present invention relates to *Trichoderma reesei* strains that have increased levels of expression of the bgl1 gene resulting in enhanced extracellular  $\beta$ -glucosidase protein levels that can be used in conjunction with other compositions to produce a cellulase product having increased cellulolytic capacity.

## 15 2. State of the Art.

[0002] Cellulases are known in the art as enzymes that hydrolyze cellulose ( $\beta$ -1,4-glucan linkages), thereby resulting in the formation of glucose, cellobiose, cellooligosaccharides, and the like. As noted by Wood et al., "Methods in Enzymology", 160, 25, pages 234 et seq. (1988) and elsewhere, cellulase produced by a given microorganism is comprised of several different enzyme classes including those identified as exocellobiohydrolases (EC 3.2.1.91) ("CBH"), endoglucanases (EC 3.2.1.4) ("EG"),  $\beta$ -glucosidases (EC 3.2.1.21) ("BG"). Moreover, the fungal classifications of CBH, EG and BG can be further expanded to include multiple components within each classification. For example, multiple CBHs and EGs have been isolated from a variety of bacterial and fungal sources including *Trichoderma reesei* which contains 2 CBHs, i.e., CBH I and CBH II, and at least 3 EGs, i.e., EG I, EG II, and EG III components.

[0003] The complete cellulase system comprising components from each of the CBH, EG, and BG classifications is required to efficiently convert crystalline forms of cellulose to glucose. Isolated components are far less effective, if at all, in hydrolyzing crystalline cellulose. Moreover, a synergistic relationship is observed between the cellulase components particularly if they are of different classifications. That is to say, the effectiveness of the complete cellulase system is significantly greater than the sum of the contributions from the isolated components of the same classification. In this regard, it is known in the art that the EG components and CBH components synergistically interact to more efficiently degrade cellulose. See, for example, Wood, Biochem. Soc. Trans., 13, pp. 407-410 (1985).

[0004] The substrate specificity and mode of action of the different cellulase components varies with classification, which may account for the synergy of the combined components. For example, the current accepted mode of cellulase action is that endoglucanase components hydrolyze internal  $\beta$ -1,4-glucosidic bonds, particularly, in regions of low crystallinity of the cellulose and exo-cellobiohydrolase components hydrolyze cellobiose from the non-reducing end of cellulose. The action of endoglucanase components greatly facilitates the action of exo-cellobiohydrolases by creating new chain ends which are recognized by exo-cellobiohydrolase components.

[0005]  $\beta$ -Glucosidases are essential components of the cellulase system and are important in the complete enzymatic breakdown of cellulose to glucose. The  $\beta$ -glucosidase enzymes can catalyze the hydrolysis of alkyl and/or aryl  $\beta$ -D-glucosides such as methyl  $\beta$ -D-glucoside and p-nitrophenyl glucoside, as well as glycosides containing only carbohydrate residues, such as cellobiose. The catalysis of cellobiose by  $\beta$ -glucosidase is important because it produces glucose for the microorganism and further because the accumulation of cellobiose inhibits cellobiohydrolases and endoglucanases thus reducing the rate of hydrolysis of cellulose to glucose.

[0006] Since  $\beta$ -glucosidases can catalyze the hydrolysis of a number of different substrates, the use of this enzyme in a variety of different applications is possible. For instance, some  $\beta$ -glucosidases can be used to liberate aroma in fruit by catalyzing various glucosides present therein. Similarly, some  $\beta$ -glucosidases can hydrolyze grape monoterpenyl  $\beta$ -glucosidase which upon hydrolysis, represents an important potential source of aroma to wine as described by Günata et al, "Hydrolysis of Grape Monoterpenyl  $\beta$ -D-Glucosides by Various  $\beta$ -Glucosidases", J. Agric. Food Chem., Vol. 38, pp. 1232-1236 (1990).

[0007] Furthermore, cellulases can be used in conjunction with yeasts to degrade biomass to ethanol wherein the cellulose degrades cellobiose to glucose that yeasts can further ferment into ethanol. This production of ethanol from readily available sources of cellulose can provide a stable, renewable fuel source. The use of ethanol as a fuel has many advantages compared to petroleum fuel products such as a reduction in urban air pollution, smog, and ozone levels, thus enhancing the environment. Moreover, ethanol as a fuel source would reduce the reliance on foreign oil imports and petrochemical supplies.

[0008] But the major rate limiting step to ethanol production from biomass is the insufficient amount of  $\beta$ -glucosidase in the system to efficiently convert cellobiose to glucose. Therefore, a cellulase composition that contains an enhanced amount of  $\beta$ -glucosidase would be useful in ethanol production.

[0009] Contrarily, in some cases, it is desirable to produce a cellulase composition which is deficient in, and preferably free of  $\beta$ -glucosidase. Such compositions would be advantageous in the production of cellobiose and other celooligosaccharides.

[0010]  $\beta$ -glucosidases are present in a variety of prokaryotic organisms, as well as eukaryotic organisms. The gene encoding  $\beta$ -glucosidase has been cloned from several prokaryotic organisms and the gene is able to direct the synthesis of detectable amounts of protein in *E. coli* without requiring extensive genetic engineering, although, in some cases, coupling with a promotor provided by the vector is required. However,  $\beta$ -glucosidases are not produced by such organisms in commercially feasible amounts.

[0011] Furthermore, such prokaryotic genes often cannot be expressed and detected after transformation of the eukaryotic host. Thus, in order to use fungal strains, fungal genes would have to be cloned using methods described herein or by detection with the *T. reesei* *bgl1* gene by nucleic acid hybridization.

[0012] The contribution and biochemistry of the  $\beta$ -glucosidase component in cellulose hydrolysis is complicated by the apparent multiplicity of enzyme forms associated with *T. reesei* and other fungal sources (Enari et al, "Purification of *Trichoderma reesei* and *Aspergillus niger*  $\beta$ -glucosidase", *J. Appl. Biochem.*, Vol. 3, pp. 157-163 (1981); Umile et al, "A constitutive, plasma membrane bound  $\beta$ -glucosidase in *Trichoderma reesei*", *FEMS Microbiology Letters*, Vol. 34, pp. 291-295 (1986); Jackson et al, "Purification and partial characterization of an extracellular  $\beta$ -glucosidase of *Trichoderma reesei* using cathodic run, polyacrylamide gel electrophoresis", *Biotechnol. Bioeng.*, Vol. 32, pp. 903-909 (1988)). These and many other authors report  $\beta$ -glucosidase enzymes ranging in size from 70-80 Kd and in pI from 7.5-8.5. More recent data suggests that the extracellular and cell wall associated forms of  $\beta$ -glucosidase are the same enzyme (Hofer et al, "A monoclonal antibody against the alkaline extracellular  $\beta$ -glucosidase from *Trichoderma reesei*: reactivity with other *Trichoderma*  $\beta$ -glucosidases", *Biochim. Biophys. Acta*, Vol. 992, pp. 298-306 (1989); Messner and Kubicek, "Evidence for a single, specific  $\beta$ -glucosidase in cell walls from *Trichoderma reesei* QM9414", *Enzyme Microb. Technol.*, Vol. 12, pp. 685-690 (1990)) and that the variation in size and pI is a result of post translational modification and heterogeneous methods of enzyme purification. It is unknown whether the intracellular  $\beta$ -glucosidase species with a pI of 4.4 and an apparent molecular weight of 98,000 is a novel  $\beta$ -glucosidase (Inglin et al, "Partial purification and characterization of a new intracellular  $\beta$ -glucosidase of *Trichoderma reesei*", *Biochem. J.*, Vol. 185, pp. 515-519 (1980)) or a proteolytic fragment of the alkaline extracellular  $\beta$ -glucosidase associated with another protein (Hofer et al, *supra*).

[0013] Since a major part of the detectable  $\beta$ -glucosidase activity remains bound to the cell wall (Kubicek, "Release of carboxymethylcellulase and  $\beta$ -glucosidase from cell walls of *Trichoderma reesei*", *Eur. J. Appl. Biotechnol.*, Vol. 13, pp. 226-231 (1981); Messner and Kubicek, *supra*; Messner et al, "Isolation of a  $\beta$ -glucosidase binding and activating polysaccharide from cell walls of *Trichoderma reesei*", *Arch. Microbiol.*, Vol. 154, pp. 150-155 (1990)), commercial preparations of cellulase are thought to be reduced in their ability to produce glucose because of relatively low concentrations of  $\beta$ -glucosidase in the purified cellulase preparation.

[0014] To overcome the problem of  $\beta$ -glucosidase being rate limiting in the production of glucose from cellulose using cellulase produced by a filamentous fungi, the art discloses supplementation of the cellulolytic system of *Trichoderma reesei* with the  $\beta$ -glucosidase of *Aspergillus* and the results indicate an increase in rate of saccharification of cellulose to glucose. Duff, *Biotechnol Letters*, 7, 185 (1985). Culturing conditions of the fungi have also been altered to increase  $\beta$ -glucosidase activity in *Trichoderma reesei* as illustrated in Sternberg et al, *Can. J. Microbiol.*, 23, 139 (1977) and Tangnu et al, *Biotechnol. Bioeng.*, 23, 1837 (1981), and mutant strains obtained by ultraviolet mutation have been reported to enhance the production of  $\beta$ -glucosidase in *Trichoderma reesei*. Although these aforementioned methods increase the amount of  $\beta$ -glucosidase in *Trichoderma reesei*, the methods lack practicality and, in many instances, are not commercially feasible.

[0015] A genetically engineered strain of *Trichoderma reesei* or other filamentous fungi that produces an increased amount of  $\beta$ -glucosidase would be ideal, not only to produce an efficient cellulase system, but to further use the increased levels of expression of the *bgl1* gene to produce a cellulase product that has increased cellulolytic capacity. Such a strain can be feasibly produced using transformation.

[0016] But, in order to transform mutant strains of *Trichoderma reesei* or other filamentous fungi, the amino acid sequence of the *bgl1* gene of *Trichoderma reesei* or the other filamentous fungi must be first characterized so that the *bgl1* gene can be cloned and introduced into mutant strains of *Trichoderma reesei* or other filamentous fungi.

[0017] Additionally, once the *bgl1* gene has been identified, information within linear fragments of the *bgl1* gene can be used to prepare strains of *Trichoderma reesei* and other filamentous fungi which produce cellulase compositions free of  $\beta$ -glucosidase.

[0018] Accordingly, this invention is directed, in part, to the characterization of the *bgl1* gene that encodes for extracellular or cell wall bound  $\beta$ -glucosidase from *Trichoderma reesei* and other filamentous fungi. This invention is further directed to the cloning of the *bgl1* gene into a plasmid vector that can be used in the transformation process, and to introduce the *bgl1* gene into the *Trichoderma reesei* or other filamentous fungi genome in multiple copies, thereby generating transformed strains which produce a cellulase composition having a significant increase in  $\beta$ -glucosidase activity. Moreover, cellulase compositions that contain increased cellulolytic capacity are also disclosed.

[0019] This invention is further directed, in part, to altered copies of the bgl1 gene which may change the properties of the enzyme and which can be reintroduced back into the Trichoderma reesei or other filamentous fungi genome.

## SUMMARY OF THE INVENTION

[0020] The amino acid sequence of the extracellular or cell wall bound  $\beta$ -glucosidase protein from Trichoderma reesei has now been obtained in sufficient detail to enable the bgl1 gene to be cloned into a suitable plasmid vector. The plasmid vector can then be used to transform strains of filamentous fungi to produce transformants which have multiple copies of the bgl1 gene introduced therein.

[0021] Accordingly, in one of its process aspects, the present invention relates to a process for expressing enhanced extracellular  $\beta$ -glucosidase in a filamentous fungus comprising expressing a fungal DNA sequence encoding enhanced  $\beta$ -glucosidase in a recombinant host microorganism, said recombinant host microorganism being a filamentous fungus transformed with an expression vector containing said DNA sequence.

[0022] In yet another process aspect, the present invention relates to a process for expressing an altered extracellular  $\beta$ -glucosidase in a filamentous fungus.

[0023] In another aspect, the present invention is directed to the amino acid sequence of extracellular  $\beta$ -glucosidase from Trichoderma reesei.

[0024] In yet another aspect, the present invention is directed to use of a nucleic acid fragment comprising the entire or partial nucleotide sequence of the T. reesei extracellular  $\beta$ -glucosidase gene as a probe to identify and clone out the equivalent bgl1 gene from other  $\beta$ -glucosidic filamentous fungi.

[0025] In one of its composition aspects, the present invention is directed to novel and useful transformants of Trichoderma reesei, which can be used to produce fungal cellulase compositions, especially fungal cellulase compositions enriched in  $\beta$ -glucosidase or deleted of  $\beta$ -glucosidase. Also contemplated in the present invention is the alteration of the bgl1 gene and the introduction of the altered bgl1 gene into T. reesei to produce transformants which can also be used to produce altered fungal cellulase compositions.

[0026] In another composition aspect, the present invention is directed to fungal cellulase compositions prepared via the transformed Trichoderma reesei strains.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0027]

Fig. 1 is the nucleotide sequence and deduced primary amino acid structure of the entire T. reesei bgl1 gene.

Fig. 2 is a schematic representation of the vector pSAS $\beta$ -glu.

Fig. 3A is a figurative representation of the vector pSAS $\Delta\beta$ Glu bal pyr ( $\Delta$ 36).

Fig. 3B is a figurative representation of the vector pUC $\Delta\beta$ -Glu A/R pyr ( $\Delta$ 12).

Fig. 4 represents a Northern blot of total RNA isolated from the transformed strains of Trichoderma reesei following induction with sophorose using the probes of cbh2 and a 700 bp fragment of bgl1 cDNA.

Fig. 5A represents an autoradiograph of a Southern blot of T. reesei DNA illustrating the presence of  $\beta$ -glucosidase gene in wild type T. reesei (RL-P37) compared to strains of T. reesei genetically modified so as to not include the  $\beta$ -glucosidase gene ( $\Delta$ 12 and  $\Delta$ 36).

Fig. 5B represents an autoradiograph of a Northern blot of T. reesei RNA illustrating the expression of  $\beta$ -glucosidase gene in wild type T. reesei (RL-P37) compared to strains of T. reesei genetically modified so as to not include the  $\beta$ -glucosidase gene ( $\Delta$ 12 and  $\Delta$ 36).

Fig. 5C represents an analysis of the proteins expressed by P37 (wild type),  $\Delta$ 12, and  $\Delta$ 36 strains of Trichoderma reesei and illustrates the absence of  $\beta$ -glucosidase in the proteins expressed by  $\Delta$ 12 and  $\Delta$ 36 strains Trichoderma reesei.

Fig. 6 represents an autoradiograph of Hind III digested genomic DNA from a T. reesei overproducing strain (lane 9) and transformants of pSAS $\beta$ -Glu (lanes 1-8), blotted and probed with the 700 bp  $\beta$ -Glu probe.

Fig. 7 represents a curve illustrating Avicel hydrolysis using the dosage, substrate:enzyme of 80:1 from an enriched recombinant  $\beta$ -glucosidase composition produced by the present invention.

Fig. 8 represents a curve illustrating PSC hydrolysis using the dosage, substrate:enzyme of 300:1 from an enriched recombinant  $\beta$ -glucosidase composition produced by the present invention.

Fig. 9 represents a curve illustrating the rate of hydrolysis of a cellulosic diaper derived fibers using an enriched recombinant  $\beta$ -glucosidase composition produced by the present invention.

Figs. 10A and 10B are autoradiographs of Aspergillus nidulans, Neurospora crassa, Humicola grisea genomic DNA digested with Hind III and Eco RI, blotted and probed with a DNA fragment containing the bgl1 gene of Trichoderma reesei.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS OF THE INVENTION

[0028] As used herein, the term "enhanced extracellular  $\beta$ -glucosidase" or "enhanced  $\beta$ -glucosidase" means that at least one additional copy of a gene encoding for extracellular  $\beta$ -glucosidase has been introduced into the genome.

5 [0029] The term "altered  $\beta$ -glucosidase" or "altered  $\beta$ -glucosidase gene" means that the amino acid sequence of the expressed protein has been altered by removing, adding, and/or manipulating the nucleic acid sequence of the gene or the amino acid sequence of the protein.

[0030] The term "by recombinant means" denotes that a microorganism has been transformed with a DNA molecule created in a test-tube by ligating together pieces of DNA that are not normally contiguous.

10 [0031] The term "cellulase free of extracellular  $\beta$ -glucosidase" refers to a cellulase composition which does not contain functional extracellular  $\beta$ -glucosidase enzyme. Such compositions are preferably prepared by culturing a filamentous fungi wherein the  $\beta$ -glucosidase gene has been either deleted or disrupted. Preferably, these compositions are prepared by culturing a filamentous fungi wherein the  $\beta$ -glucosidase gene has been deleted.

[0032] The term "filamentous fungi" means any and all art recognized filamentous fungi.

15 [0033] The term " $\beta$ -glucosidic filamentous fungi" refers to those filamentous fungi which produce a cellulase composition containing  $\beta$ -glucosidase.

[0034] The term "cellooligosaccharide" refers to those oligosaccharide groups containing from 2-8 glucose units having  $\beta$ -1,4 linkages. Such cellooligosaccharides include cellobiose (diglucose having a  $\beta$ -1,4- linkage) and are preferably derived from cellulose.

20 [0035] More specifically, the present invention relates to the isolation and characterization of the bgl1 gene coding for the extracellular or cell wall bound protein from Trichoderma reesei (sometimes referred to as "T. reesei") and the specific nucleotide and amino acid sequence of this gene. The bgl1 gene is cloned into plasmid vectors, which are further used to produce transformed strains of T. reesei and other filamentous fungi having extra copies of the bgl1 gene inserted therein. These transformants are then used to produce cellulase compositions having increased  $\beta$ -glucosidase activity and thus enhanced cellulolytic degradation.

25 [0036] Also contemplated by the present invention is the manipulation of the amino acid sequence in the bgl1 gene itself. Alteration of the active sites on this enzyme may lead to a variety of different changes in catalytic conversion. For example, since  $\beta$ -glucosidase has both hydrolase and transferase activity, alteration of the amino acid sequence may result in the removal of hydrolase activity and an increase in transferase activity and, thus, facilitate the synthesis of  $\beta$  1-4 oligo-dextrins. Moreover, manipulation of the amino acid sequence of  $\beta$ -glucosidase may result in further changes in the system, such as different pH optima, different temperature optima, altered catalytic turn over rate ( $V_{max}$ ), altered affinity ( $K_m$ ) for cellobiose leading to an increased affinity for cellobiose or a decreased affinity for cellobiose resulting in a slower or zero rate of reaction, altered product inhibition profile such that lower or higher levels of glucose will inhibit  $\beta$ -glucosidase activity, and the like.

30 [0037] Moreover, a nucleic acid fragment containing the entire nucleotide sequence of the extracellular  $\beta$ -glucosidase gene in T. reesei or a portion thereof can also be labeled and used as a probe to identify and clone out the equivalent bgl1 gene in other filamentous fungi.

35 [0038] Generally, the present invention involves the isolation of the bgl1 gene from T. reesei by identifying a 700 bp cDNA fragment of the gene which is then used as a probe to identify a single T. reesei fragment containing the bgl1 gene which was subsequently cloned. Because of the species homology of the bgl1 gene, a probe employing a fragment of the bgl1 gene of T. reesei can be employed to identify the bgl1 gene in other cellulolytic microorganisms and, it is understood that the following description for T. reesei could also be applied to other  $\beta$ -glucosidic filamentous fungi.

40 [0039] In the case of T. reesei, this 6.0 kb fragment is then cloned into a pUC plasmid and a series of mapping experiments are performed to confirm that the entire bgl1 gene is contained in this fragment. The nucleotide sequence is then determined on both strands and the position of two introns can be confirmed by sequence analysis of bgl1 cDNA subclones spanning the intron/exon boundaries. After isolation of the bgl1 gene, additional bgl1 gene copies are then introduced into T. reesei or other filamentous fungal strains to increase the expression of  $\beta$ -glucosidase.

45 [0040] The isolation of the bgl1 gene from T. reesei involves the purification of extracellular  $\beta$ -glucosidase, chemical and proteolytic degradation of this protein, isolation and determination of the sequence of the proteolytic fragments and design of synthetic oligomer DNA probes using the protein sequence. The oligomeric probes are then further used to identify a 700 bp  $\beta$ -glucosidase cDNA fragment which can be labeled and employed to later identify a fragment that contains the entire bgl1 gene within the fragment from digested genomic DNA from T. reesei.

50 [0041] To identify a feasible cDNA fragment that can be used as a probe for future analysis, total RNA is first isolated from T. reesei mycelia and polyadenylated RNA isolated therefrom. The polyadenylated RNA is then used to produce a cDNA pool which is then amplified using specific oligonucleotide primers that amplify only the specific cDNA fragment encoding the T. reesei bgl1 gene.

55 [0042] More specifically, total RNA is first isolated from a starting strain of T. reesei. The starting strain employed in the present invention can be any T. reesei cellulase overproduction strain that is known in the art. This cellulase pro-

ducing strain is generally developed by ordinary mutagenesis and selection methods known in the art from any *T. reesei* strain. Confirmation that the selected strain overproduces cellulases can be performed by using known analysis methods. A preferred strain is RLP37 which is readily accessible.

5 [0043] A mycelial inoculum from the *T. reesei* over production strain, grown in an appropriate growth medium, is added to a basal medium and incubated for a period of between 50-65 hours at a temperature between 25°C to 32°C, preferably 30°C. Fresh basal medium can be replaced during this incubation period. The culture medium is then centrifuged, and the mycelia is isolated therefrom and washed. The mycelia is then resuspended in a buffer to permit growth thereof and 1 mM sophorose (a  $\beta$ ,1-2 dimer of glucose) is added to the mycelia to induce the production of cellulase enzymes. The mycelia preparation is then incubated for an additional time period, preferably 18 hours at 10 30°C prior to harvesting.

[0044] Total RNA can be isolated from the mycelia preparation by a variety of methods known in the art, such as proteinase K lysis, followed by phenol:chloroform extraction, guanidinium isothiocyanate extraction, followed by cesium chloride gradients, guanidine hydrochloride and organic solvent extraction, and the like. It is preferable to isolate total RNA via the procedure described by Timberlake et al in "Organization of a Gene Cluster Expressed Specifically in the 15 Asexual Spores of *A. nidulans*," *Cell*, 26, pp. 29-37 (1981). The mycelia is isolated from the culture medium via filtration. Then the RNA is extracted from the mycelia by the addition of an extraction buffer, TE-saturated phenol and chloroform. The aqueous phase is removed and the organic phase is reextracted with the extraction buffer alone by heating the extraction mixture in a water bath at a temperature between about 60°C to 80°C, preferably 68°C to release the RNA trapped in polysomes and at the interface. All of the extracted aqueous phases are then pooled, centrifuged and re-extracted with phenol-chloroform until there is no longer any protein at the interface. The RNA is further precipitated 20 with 0.1 volume of 3 M sodium acetate and 2 volumes of 95% ethanol and pelleted via centrifugation before it is resuspended in DEP-water containing an RNase inhibitor.

[0045] The total RNA is then fractionated on 1% formaldehyde-agarose gels, blotted to Nytran™ membranes, and probed using a fragment of the *T. reesei cbh2* gene to determine whether the genes encoding the enzymes of the 25 cellulase system in the *T. reesei* preparation are indeed induced by addition of the sophorose. Basically, the probe used in the present invention is derived from a CBH II clone produced by methods known in the art. For more specific detail of how the clone was produced see Chen et al, "Nucleotide Sequence and Deduced Primary Structure of Cellobiohydrolase II from *Trichoderma reesei*," *Bio/Technology*, Vol. 5 (March 1987). Site directed mutagenesis was performed on the CBH II clone and a *Bgl* II site was placed at the exact 5' end of the open reading frame and a *Nhe* I site at the 30 exact 3' end. The *Bgl* II and *Nhe* I restriction fragment containing CBH II coding sequence was further cloned into a pUC218 phagemid. The CBH II gene was further cut and gel isolated prior to adding a label.

[0046] The results of the Northern blot of *T. reesei* RNA probed with the *cbh2* probe indicated that the level of *cbh2* specific mRNA reached a peak at 14-18 hours post induction. From this data it can be inferred that the entire cellulase complex including  $\beta$ -glucosidase is induced at this time. The total RNA from 14, 18 and 22 hours is then pooled.

35 [0047] After pooling the specific fractions of total RNA, polyadenylated mRNA is further isolated from the total RNA. Postranscriptional polyadenylation is a common feature of the biogenesis of most eukaryotic mRNAs. The newly synthesized mRNAs have long poly(A) tracts which tend to shorten as mRNAs age. The newly synthesized polyadenylated mRNA is further isolated from total RNA by methods known in the art. These methods include the use of oligo(dT)-cellulose, poly(U) Sepharose, adsorption to and elution from poly(U) filters or nitrocellulose membrane filters, and the 40 like. It is preferable to use oligo(dT) cellulose chromatography in isolating mRNA following the procedure described by Sambrook et al, *Molecular Cloning, A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press (1989). More specifically, fractions of total RNA are run through the chromatographic resin, and mRNA is eluted therefrom with an elution buffer. The RNA which binds to the column is enriched for RNAs containing poly(A) tails and, therefore, eliminates contaminants, such as rRNA and partially degraded mRNAs. It is important that the purification be carried 45 out successfully such that when cDNA is synthesized from the mRNA, higher yields of mRNA copies and less spurious copying of non-messenger RNAs occurs.

[0048] Total RNA and polyadenylated RNA from the preparations were further fractionated on 1% formaldehyde gels, blotted to Nytran<sup>R</sup> membranes and analyzed to confirm that the enzymes in the cellulase complex were being induced as polyadenylated mRNA.

50 [0049] After isolating polyadenylated mRNA from total RNA, complementary DNA or cDNA is synthesized therefrom. The first strand of cDNA is synthesized using the enzyme RNA-dependent DNA polymerase (reverse transcriptase) to catalyze the reaction. Avian reverse transcriptase which is purified from the particles of an avian retrovirus or murine reverse transcriptase, which is isolated from a strain of *E. coli* that expresses a cloned copy of the reverse transcriptase gene of the Moloney murine leukemia virus can be used in the present invention. However, it is preferable to use the 55 Moloney murine leukemia virus (M-MLV) reverse transcriptase to synthesize first strand cDNA from the polyadenylated mRNA population. The amount of cloned M-MLV reverse transcriptase required may vary depending on the amount of polyadenylated mRNA used in the synthesis reaction. Usually, about 200 U/ $\mu$ l of the reverse transcriptase is used per 2 to 10  $\mu$ g of mRNA per reaction.



[0050] Also present in the synthesis mixture is a primer to initiate synthesis of DNA. For cloning of cDNAs, any primer can be used, but it is preferable to use oligo(dT) containing 12-18 nucleotides in length, which binds to the poly(A) tract at the 3' terminus of eukaryotic cellular mRNA molecules. The primer is added to the reaction mixture in large molar excess so that each molecule of mRNA binds several molecules of oligo(dT)<sub>12-18</sub>. It is preferable to use about 12.5 µg of primer having a concentration of 0.5 mg/ml.

[0051] Besides the enzyme and primer, a buffer and dNTP mix containing dATP, dCTP, dGTP, and dTTP at a final concentration of 500 µM each usually completes the reaction cocktail. Any buffer can be used in the present invention for first strand cDNA synthesis that is compatible with this synthesis. It is preferable to use a buffering system consisting of 250 mM Tris-HCl (pH 8.3), 375 mM KCl, 15 mM MgCl<sub>2</sub>, and 50 mM dithiothreitol. Generally, about 500 µl of buffer completes the synthesis solution.

[0052] After the first strand is synthesized, the second strand of cDNA may be synthesized by a variety of methods known in the art, such as hairpin-primed synthesis by denaturing the cDNA:mRNA complex, adding the Klenow fragment of *E. coli* DNA polymerase or reverse transcriptase, and then digesting the hairpin loop with nuclease S1 to obtain a double-stranded cDNA molecule, the Okayama and Berg method, the Gubler and Hoffman method, and the like. The Okayama and Berg method uses *E. coli* RNase H to randomly nick the mRNA, and the RNA is replaced in the nick translation reaction by catalysis with *E. coli* DNA polymerase I. In the Okayama and Berg method, mRNA is used to prime DNA synthesis by the *E. coli* DNA polymerase I.

[0053] The preferred method to synthesize the second strand of cDNA is a modified method of the Gubler and Hoffman procedure. This procedure uses *E. coli* RNase H, DNA Polymerase I, and DNA Ligase to form the second strand. Actually, two different methods of proceeding with the second strand synthesis can be used in the present invention. The first procedure uses RNase H to attack the RNA:DNA hybrid in a random fashion, producing nicks in addition to those produced by reverse transcriptase. If too many nicks are introduced into the RNA at the 5' end of the message before second strand synthesis commences, fragments may be produced that are too short to remain hybridized; thus, they will not be able to serve as primers. In addition, the 5'-most RNA oligomer which primes second strand DNA synthesis will continue to be degraded until only two ribonucleotides remain at the 5' end of the second strand DNA. These are substrates for the polymerase I RNase H activity, and the remaining nucleotides will be removed. This leaves the 3' end of the first strand cDNA single stranded, making it a substrate for the 3' exonuclease activity of Polymerase I. The result is a population of cDNAs, which are blunt-ended.

[0054] An alternative method relies on M-MLV reverse transcriptase to produce nicks 10 to 20 bases from the 5' end of the RNA in the hybrid. DNA polymerase I is then used for synthesis. Generally, about 500 units at a concentration of 10 U/µl of DNA polymerase I is used. After second strand synthesis, RNase H is added after removal of the DNA polymerase I to produce a duplex, which is entirely DNA, except for the surviving capped RNA 5' oligonucleotide.

[0055] The second-strand synthesis by either procedure set forth above usually takes place in the presence of a buffer and dNTP mix. Any buffering system that is known in the art for second strand cDNA synthesis can be used; however, it is preferable to use a buffering system containing 188 mM Tris-HCl, pH 8.3, 906 mM KCl, 100 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 46 mM MgCl<sub>2</sub>, 37.5 mM dithiothreitol, and 1.5 mM NAD. The dNTP mix preferably contains 10 mM dATP, 10 mM dCTP, 10 mM dGTP, and 10 mM dTTP.

[0056] The second strand synthesis is carried out under known procedures set forth in the art. The preferred methods and reagents used to synthesize cDNA in the present invention are the BRL cDNA Synthesis System<sup>R</sup> (Bethesda Research Laboratories, Gaithersburg, Maryland) and the Librarian System (Invitrogen, San Diego, CA).

[0057] At this point a pool of cDNAs, a small portion of which code for the *bgl1* gene, is present after second strand synthesis. Since amplification of only the specific *bgl1* gene fragment in the cDNA pool is crucial for the isolation of the β-glucosidase gene, specific primers were designed to amplify the cDNA fragment encoding the *T. reesei bgl1* gene in the polymerase chain reaction (PCR). The primers used are degenerate primers designed to hybridize to the cDNA of the *bgl1* gene encoding the N-terminus and an internal CNBr fragment.

[0058] In general, it is difficult to isolate the *bgl1* gene because the amino acid sequence of the protein does not contain sufficient amino acids which are coded for by unique nucleic acid triplets and thus any oligonucleotide used would be too degenerate to specifically amplify the *bgl1* gene in the PCR reaction. However, in this invention, primers were designed by examining the amino acids of the region targeted for amplification of mature β-glucosidase and choosing regions, which will require a reduced degree of degeneracy in the genetic code. Codon bias in *T. reesei* for various other cellulase genes such as *cbh1*, *cbh2*, *egl1*, and the like was also taken into account when designing the oligonucleotide primers. More specifically, codon bias is based on various genes in the strain *T. reesei* which display a preferred nucleotide triplet encoding different amino acids. By analyzing this codon bias one can determine that a particular nucleotide sequence coding for an amino acid would be preferred. For example, the *cbh1*, *cbh2* and *egl1* genes from *T. reesei* prefer the CCU coding for the amino acid proline. Thus, when designing an oligonucleotide probe, the CUG sequence would be the preferred choice for leucine, rather than the other triplets (CUU, CUC, CUA, UUA and CUG) which code for leucine.

[0059] Furthermore, after selection of an N-terminal region and an internal region as primers for amplification pur-

poses, the primers were designed by inserting a non-specific base inosine into the wobble position of the primer for the 5' terminus and using a pool of sixteen variable primer sequences for the internal primer. Basically, the creation of degenerate primers is described by Compton in "Degenerate Primers For DNA Amplification" and Lee et al in "cDNA Cloning Using Degenerate Primers" in PCR Protocols: A Guide to Methods and Applications, published by Academic Press (1990).

[0060] Using the primers described above, the cDNA sequences encoding the amino terminal region of the *bgl1* gene is then selectively amplified using PCR. The amplification method consists of an initial, denaturing cycle of between about 5 to 15 minutes at 95°C, followed by a 1-7 minutes annealing step at a temperature between 35°C and 55°C and preferably between 45°C and 55°C and a 5-15 minutes polymerization cycle at 65°C. It is preferable, however, to use a 10 minute initial denaturing cycle, followed by 2 minutes of annealing at 50°C and a 10 minute, and preferably a 30 minute polymerization cycle at the aforescribed temperatures.

[0061] The amplified fragment is then identified via gel electrophoresis as a 700 bp cDNA segment. The amplified pool of cDNAs is then further fractionated on a polyacrylamide gel to obtain a more purified 700 bp cDNA fragment for cloning purposes. After elution of the 700 bp fragments from the gel, the 700 bp cDNA fragments are then cloned into phagemid vectors. Any cloning vector can be used to clone the cDNA *bgl1* gene fragments, such as pUC18, pUC19, pUC118, pUC119, pBR322, pEMBL, pRSA101, pBluescript, and the like. However, it is preferable to use the cloning vectors pUC218 and pUC219, which are derived from pUC18 and pUC19 by insertion of the intergenic region of M13. The cloning vectors with the cDNA fragments containing the *bgl1* gene are then used to transform *E. coli* strain JM101. After transformation, positive colonies containing the *bgl1* gene were identified and DNA isolated therefrom using chloroform:phenol extraction and ethanol precipitation methods.

[0062] The nucleotide sequence of the subcloned cDNA 700 bp fragment is then determined by the dideoxy chain termination method described by Sanger et al using a Sequenase<sup>R</sup> reagent kit provided by U.S. Biochemicals.

[0063] From this nucleotide sequence it was determined that the subcloned 700 bp cDNA segment contained an open reading frame encoding 150 amino acids that overlapped a number of other sequenced peptides that were obtained following CNBr and proteolytic degradation of purified *T. reesei*  $\beta$ -glucosidase. Thus, it was confirmed that the cloned sequences encoded for the extracellular *T. reesei*  $\beta$ -glucosidase protein.

[0064] The cloning of the genomic version of the entire  $\beta$ -glucosidase gene was then undertaken by labelling the 700 bp *bgl1* cDNA fragment with <sup>32</sup>P using the methods to label oligonucleotides described by Sambrook et al, *supra*. This probe is used to identify a 6.0 kb band on a Southern blot of *Hind* III digested genomic DNA from *T. reesei*.

[0065] The genomic DNA from *T. reesei* is prepared for Southern blot analysis by deproteinizing the genomic DNA, followed by treatment with ribonuclease A. The prepared genomic DNA is then cut with one of a variety of restriction enzymes such as *Eco* RI, *Hind* III and the like, run on a gel, Southern blotted and hybridized with the 700 bp cDNA labelled fragment of the *bgl1* gene. From this analysis, it was determined that *Hind* III was the restriction enzyme of choice that can be used to clone the *bgl1* gene.

[0066] *Hind* III is then added to genomic DNA from the strain *T. reesei* and DNA is extracted therefrom. A sample from this digestion is run on an agarose gel and fractionated electrophoretically. The gel is then Southern blotted and probed with the 700 bp cDNA probe. A 6.0 kb band was then identified on the Southern blot of *Hind* III digested genomic DNA. The remaining *Hind* III digested genomic DNA was then subjected to preparative gel electrophoresis and DNA ranging in size from about 5.0 kb to 7.0 kb was eluted therefrom and cloned into a phagemid vector and used to transform *E. coli* JM101 to create a library. Any phagemid vector can be used such as those described above, however it is preferable to use pUC218. The colonies that resulted from the transformation were then subjected to colony hybridization using the 700 bp cDNA fragment as a probe to identify those colonies that contained the cloned genomic DNA coding for *bgl1*. The positive colonies from the transformation are then picked and the DNA isolated therefrom by methods known in the art.

[0067] The isolated DNA from such a positive colony is then digested with various restriction enzymes, both singly and in various combinations, and subjected to agarose gel electrophoresis. The resultant banding pattern is then used to construct a restriction map of the cloned 6.0 kb genomic DNA from *T. reesei*. Enzymes used in the digestion include *Eco* RI, *Sst* I, *Kpn* I, *Sma* I, *Bam* HI, *Xho* I, *Bgl* II, *Cla* I, *Xba* I, *Sal* I, *Pst* I, *Sph* I, *Hind* III, *Bal* I, *Pvu* II and the like.

[0068] The same gel is then subject to Southern blot analysis using the same 700 bp *bgl1* cDNA as a probe to identify which genomic restriction fragments shared homology with the *bgl1* cDNA. Since the position of these homologous fragments can be determined relative to the restriction map of the 6.0 kb genomic fragment and also since the size of the  $\beta$ -glucosidase protein (74 kd) gives an estimated length of the gene as 2.1 kb (because average molecular weight of an amino acid is 105 daltons, a 74 kd protein contains on average 705 amino acids, which in turn is equal to 2,115 bp), then the mapping experiments confirmed that the entire *bgl1* gene is contained on the genomic *Hind* III clone.

[0069] *Pvu* II and *Bal* I restriction fragments ranging in size from 600 bp to 1500 bp hybridized with the 700 bp cDNA *bgl1* clone and were thus chosen for subcloning into pUC218 phagemids. The nucleotide sequence was determined using the methods of Sanger et al, described above. The *Pvu* II and *Bal* I subclones were sequenced and the overlapping sequences of the subclones aligned until a single contiguous sequence totaling 3033 bp was obtained within which

the nucleotide sequence of the bgl1 gene was determined on both strands and the position of two small introns was inferred by homology to introns of other genes of filamentous fungi. The amino acid sequence is also deduced as set forth in Figure 1.

[0070] The nucleotide sequence and deduced primary amino acid sequence of the entire T. reesei bgl1 gene is set forth in Figure 1. The predicted molecular weight of the encoded  $\beta$ -glucosidase protein is 74,341. A 31 amino acid peptide precedes the mature amino terminus of  $\beta$ -glucosidase as deduced from the amino terminal peptide sequence. Within this peptide, there are three potential signal peptidase recognition sites consisting of Ala-X-Ala.

[0071] The primary amino acid sequence of  $\beta$ -glucosidase shows 7 potential N-linked glycosylation sites at positions 208, 310, 417, and 566, which shows the consensus Asn-X-Ser/Thr-X where X is not a proline. However, sites at positions 45, 566, and 658 have a proline residue in the consensus sequence and may or may not be glycosylated.

[0072] No unusual codon bias is observed in the bgl1 gene when compared to other cellulase genes. The bgl1 coding region is interrupted by two short introns of 70 bp and 64 bp, respectively. Both introns have splice site donor, splice acceptor, and lariat branch acceptor sites that show homology to the consensus splice signals emerging from T. reesei and other filamentous fungi.

[0073] Since the bgl1 gene from the T. reesei strain is identified and can be cloned, the next step is to produce a transformant that has extra copies of the bgl1 gene.

[0074] A selectable marker must first be chosen so as to enable detection of the transformed filamentous fungus. Different selectable markers may be used including argB from A. nidulans or T. reesei, amdS from A. nidulans, pyr4 from Neurospora crassa, A. nidulans or T. reesei, and pyrG from Aspergillus niger. The selectable marker can be derived from a gene, which specifies a novel phenotype, such as the ability to utilize a metabolite that is usually not metabolized by the filamentous fungi to be transformed or the ability to resist toxic shock effects of a chemical or an antibiotic. Also contemplated within the present invention are synthetic gene markers that can be synthesized by methods known in the art. Transformants can then be selected on the basis of the selectable marker introduced therein. Because T. reesei does not contain the amdS gene, it is preferable to use the amdS gene in T. reesei as a selectable marker that encodes the enzyme acetamidase, which allows transformant cells to grow on acetamide as a nitrogen source. In the case where the bgl1 gene is deleted from T. reesei, it is preferable to use the pyrG gene as a selectable marker.

[0075] The host strain used should be mutants of the filamentous fungi which lack or have a nonfunctional gene or genes corresponding to the selectable marker chosen. For example, if the selectable marker of argB is used, then a specific arg<sup>-</sup> mutant strain is used as a recipient in the transformation procedure. Other examples of selectable markers that can be used in the present invention include the genes trp, pyr4, pyrG, trp1, oliC31, Bm1, pkIA, niaD, leu, and the like. The corresponding recipient strain must, therefore, be a mutant strain such as trp<sup>-</sup>, pyr<sup>-</sup>, leu<sup>-</sup>, and the like.

[0076] The mutant strain is derived from a starting host strain, which is any filamentous fungi strain. However, it is preferable to use a filamentous fungi over-producing mutant strain and particularly, a T. reesei overproducing strain described previously, since this strain secretes high amounts of proteins and, in particular, high amounts of cellulase enzymes. The selected mutant strain is then used in the transformation process. The preferred strain of T. reesei for use in deleting the bgl1 gene is RLP37 pyrG69, a uridine auxotroph.

[0077] The mutant strain of the selected filamentous fungi can be prepared by a number of techniques known in the art, such as the filtration enrichment technique described by Nevalainen in "Genetic improvement of enzyme production in industrially important fungal strains", Technical Research Center of Finland, Publications 26 (1985). Another technique to obtain the mutant strain is to identify the mutants under different growth medium conditions. For instance, the arg<sup>-</sup> mutants can be identified by using a series of minimal plates supplied by different intermediates in arginine biosynthesis. Another example is the production of pyr<sup>-</sup> mutant strains by subjecting the strains to fluoroorotic acid (FOA). Strains with an intact pyr4 gene grow in an uridine medium and are sensitive to fluoroorotic acid, and, therefore, it is possible to select pyr4<sup>-</sup> mutant strains by selecting for FOA resistance.

[0078] The chosen selectable marker is then cloned into a suitable plasmid. Any plasmid can be used in the present invention for the cloning of the selectable marker such as pUC18, pBR322, and the like. However, it is preferable to use pUC100. The vector is created by digesting pUC100 with the restriction enzyme SmaI, and the 5' phosphate groups are then removed by digestion with calf alkaline phosphatase. The fragment vector is then purified by gel electrophoresis followed by electroelution from the isolated gel slice. The amdS gene from A. nidulans is isolated as a 2.4 kb SstI restriction fragment following separation of the vector sequences via known procedures such as those described by Hynes et al, Mol. Cell. Biol., 3, pp. 1430-1439 (1983). The 2.4 Kb SstI amdS fragment and the 2.7 Kb pUC100 vector fragment are then ligated together, and the ligation mix is then introduced into the E. coli host strain JM101.

[0079] Any plasmid can be used in the present invention for the insertion of the bgl1 gene, but it is preferable to use the pSAS plasmid.

[0080] pSAS $\beta$ -glu is constructed by digesting pSAS with the restriction enzyme Hind III and purifying the linear fragment via gel electrophoresis and electroelution. Into this Hind III treated pSAS vector fragment is ligated the 6.0 Kb Hind III fragment of T. reesei genomic DNA that contained all of the coding region of the bgl1 gene along with th

sequences necessary for transcription and translation. Figure 2 illustrates the construction of pSAS $\beta$ -glu.

[0081] It is also possible to construct vectors that contain at least one additional copy of the bgl1 gene and to construct vectors in which the amino acid sequence of bgl1 gene has been altered by known techniques in the art such as site directed mutagenesis, PCR methods, and chemical mutation methods.

5 [0082] After a suitable vector is constructed, it is used to transform strains of filamentous fungi. Since the permeability of the cell wall in filamentous fungi (e.g., *T. reesei*) is very low, uptake of the desired DNA sequence, gene or gene fragment is at best minimal. To overcome this problem, the permeability of the cell wall can be increased or the DNA can be directly shot into the cells via a particle gun approach. In the particle gun approach, the DNA to be incorporated into the cells is coated onto micron size beads and these beads are literally shot into the cells leaving the DNA therein and leaving a hole in the cell membrane. The cell then self-repairs the cell membrane leaving the DNA incorporated therein. Besides this aforescribed method, there are a number of methods to increase the permeability of filamentous fungi cells walls in the mutant strain (i.e., lacking a functional gene corresponding to the used selectable marker) prior to the transformation process.

15 [0083] One method involves the addition of alkali or alkaline ions at high concentrations to filamentous fungi cells. Any alkali metal or alkaline earth metal ion can be used in the present invention; however, it is preferable to use either CaCl<sub>2</sub> or lithium acetate and more preferable to use lithium acetate. The concentration of the alkali or alkaline ions may vary depending on the ion used, and usually between 0.05 M to 0.4 M concentrations are used. It is preferable to use about a 0.1 M concentration.

20 [0084] Another method that can be used to induce cell wall permeability to enhance DNA uptake in filamentous fungi is to resuspend the cells in a growth medium supplemented with sorbitol and carrier calf thymus DNA. Glass beads are then added to the supplemented medium, and the mixture is vortexed at high speed for about 30 seconds. This treatment disrupts the cell walls, but may kill many of the cells.

25 [0085] Yet another method to prepare filamentous fungi for transformation involves the preparation of protoplasts. Fungal mycelium is a source of protoplasts, so that the mycelium can be isolated from the cells. The protoplast preparations are then protected by the presence of an osmotic stabilizer in the suspending medium. These stabilizers include sorbitol, mannitol, sodium chloride, magnesium sulfate, and the like. Usually, the concentration of these stabilizers varies between 0.8 M to 1.2 M. It is preferable to use about a 1.2 M solution of sorbitol in the suspension medium.

30 [0086] Uptake of the DNA into the host mutant filamentous fungi strain is dependent upon the concentration of calcium ion. Generally, between about 10 mM CaCl<sub>2</sub> and 50 mM CaCl<sub>2</sub> is used in an uptake solution. Besides the need for calcium ions in the uptake solution, other items generally included are a buffering system such as TE buffer (10 mM Tris, pH 7.4; 1 mM EDTA) or 10 mM MOPS, pH 6.0 buffer (morpholinepropane-sulfonic acid), and polyethylene glycol (PEG). The polyethylene glycol acts to fuse the cell membranes, thus permitting the contents of the mycelium to be delivered into the cytoplasm of the filamentous fungi mutant strain and the plasmid DNA is transferred to the nucleus. This fusion frequently leaves multiple copies of the plasmid DNA tandemly integrated into the host chromosome. Generally, a high concentration of PEG is used in the uptake solution. Up to 10 volumes of 25% PEG 4000 can be used in the uptake solution. However, it is preferable to add about 4 volumes in the uptake solution. Additives such as dimethyl sulfoxide, heparin spermidine, potassium chloride, and the like may also be added to the uptake solution and aid in transformation.

40 [0087] Usually a suspension containing the filamentous fungi mutant cells that have been subjected to a permeability treatment or protoplasts at a density of 10<sup>8</sup> to 10<sup>9</sup>/ml, preferably 2 x 10<sup>8</sup>/ml, are used in transformation. These protoplasts or cells are added to the uptake solution, along with the desired transformant vector containing a selectable marker and other genes of interest to form a transformation mixture.

45 [0088] The mixture is then incubated at 4°C for a period between 10 to 30 minutes. Additional PEG is then added to the uptake solution to further enhance the uptake of the desired gene or DNA sequence. The PEG may be added in volumes of up to 10 times the volume of the transformation mixture, preferably, about 9 times. After the PEG is added, the transformation mixture is then incubated at room temperature before the addition of a sorbitol and CaCl<sub>2</sub> solution. The protoplast suspension is then further added to molten aliquots of a growth medium. This growth medium contains no uridine and selectively permits the growth of transformants only. The subsequent colonies were transferred and purified on a growth medium depleted of sorbitol.

50 [0089] At this stage, stable transformants can be distinguished from unstable transformants by their faster growth rate and the formation of circular colonies with a smooth rather than ragged outline on solid culture medium. Additionally, in some cases, a further test of stability can be made by growing the transformants on solid non-selective medium, harvesting the spores from this culture medium and determining the percentage of these spores which will subsequently germinate and grow on selective medium.

55 [0090] In order to ensure that the transformation took place by the above-described methods, further analysis is performed on the transformants such as Southern blotting and autoradiography. Using the same basic procedures set forth above, the entire bgl1 gene can be deleted from a vector and transformed into filamentous fungi strains or the bgl1 gene can be altered and transformed into filamentous fungi strains.

[0091] After confirmation that the transformed strains contained at least one additional copy of the bgl1 gene or an altered bgl1 gene, the strains are further cultured under conditions permitting these transformants to propagate. The transformants can then be isolated from the culture media and used in a variety of applications which are described below. Alternatively, the transformants can be further fermented and a recombinant fungal cellulase composition can be isolated from the culture media. Since, for example, the transformants produced by the present invention can express enhanced or altered extracellular  $\beta$ -glucosidase in the fermentation medium, fungal cellulase compositions can be isolated from the medium. Usually, the isolation procedure involves centrifuging the culture or fermentation medium containing the transformants and filtering by ultrafiltration the supernatant to obtain a recombinantly produced fungal cellulase composition. Optionally, an antimicrobial agent can be further added to the composition prior to use in the variety of applications described below. Examples of microbial agents that can be added are sodium azide, sodium benzoate and the like.

[0092] Confirmation that the transformants produced by the process of the present invention had enhanced activity on cellobiose, the following experiment was performed. In this experiment 50 mg of cellobiose which was suspended in 1.0 ml of phosphate buffer (pH 5.0) and was reacted with the fermentation product produced by the transformant (65.5 mg/ml protein) using a fermentation product from a normal nonmutant T. reesei strain as a control (135.0 mg/ml protein). The results of cellobiose activity under conditions of initial rate, are set forth in Table I below:

TABLE I

Product	Protein (mg/ml)	Activity on Cellobiose $\mu$ mole glucose
		mg protein
Control	135.0	6
Product produced by the present invention	65.5	33

[0093] The results from this experiment indicate that the fermentation product produced by the transformants of the present invention has over five times the specific activity on the substrate, cellobiose, compared to a nonmutant T. reesei control strain.

[0094] Moreover, Figures 7 and 8 confirm that hydrolysis is enhanced for the substrates Avicel and PSC (note: PSC is a phosphoric acid swollen cellulose obtained by treating Avicel with phosphoric acid) using 1.0% enzyme/substrate. In the experiment, PSC or Avicel was suspended in 2 mls of 50 mM sodium acetate buffer, pH 4.8, and incubated at 40° under non-agitated conditions for up to 24 hours. Soluble reducing sugar was measured by the method of Nelson and Somogyi. From these figures it is further demonstrated that the enhanced recombinant  $\beta$ -glucosidase fermentation product produced from transformants according to the present invention, has an increased rate and extent of hydrolytic activity on the various substrates compared to the standard Cyt-123 control (on average 20% higher activity). The Cyt-123 control is the product obtained from a T. reesei cellulase over-production strain subjected to fermentation on an industrial scale.

[0095] The enriched transformants can be used in a variety of different applications. For instance, some  $\beta$ -glucosidases can be further isolated from the culture medium containing the enhanced transformants and added to grapes during wine making to enhance the potential aroma of the finished wine product. Yet another application can be to use  $\beta$ -glucosidase in fruit to enhance the aroma thereof. Alternatively, the isolated recombinant fermentation product containing enhanced  $\beta$ -glucosidase can be used directly in food additives or wine processing to enhance the flavor and aroma.

[0096] Since the rate of hydrolysis of cellulosic products is increased by using the transformants having at least one additional copy of the bgl1 gene inserted into the genome, products that contain cellulose or heteroglycans can be degraded at a faster rate and to a greater extent. Products made from cellulose such as paper, cotton, cellulosic diapers and the like can be degraded more efficiently in a landfill. Figure 9 illustrates the use of an increased  $\beta$ -glucosidase preparation isolated from the fermentation medium containing transformants having at least one additional copy of the bgl1 gene inserted into the genome compared to a non-enhanced Cyt 123 standard (defined above) on a cellulosic diaper product. This hydrolysis experiment was performed using 0.4 mg of the standard and the fermentation product per 100 mg of substrate (the cellulosic diaper). The experiment was run at 50°C over a period of five hours and the glucose concentration was measured, in duplicate, at various time intervals. This curve illustrates an increased rate of hydrolysis for the product produced by the fermentation product derived from the transformant having additional copies of bgl1, compared to the standard. It was also determined that the diaper derived fibers were about 14% insoluble in aqueous solution. Thus, the fermentation product obtained from the transformants or the transformants alone can be used in compositions to help degrade by liquefaction a variety of cellulose products that add to overcrowded landfills.

[0097] Simultaneous saccharification and fermentation is a process whereby cellulose present in biomass is converted to glucose and, at the same time and in the same reactor, yeast strains convert the glucose into ethanol. Y ast

strains that are known for use in this type of process include B. clausenii, S. cerevisiae, Cellulolyticus acidothermophilum, C. brassicae, C. lustinaniae, S. uvarum, Schizosaccharomyces pombe and the like. Ethanol from this process can be further used as an octane enhancer or directly as a fuel in lieu of gasoline which is advantageous because ethanol as a fuel source is more environmentally friendly than petroleum derived products. It is known that the use of ethanol will improve air quality and possibly reduce local ozone levels and smog. Moreover, utilization of ethanol in lieu of gasoline can be of strategic importance in buffering the impact of sudden shifts in non-renewable energy and petro-chemical supplies.

[0098] Ethanol can be produced via saccharification and fermentation processes from cellulosic biomass such as trees, herbaceous plants, municipal solid waste and agricultural and forestry residues. However, one major problem encountered in this process is the lack of  $\beta$ -glucosidase in the system to convert cellobiose to glucose. It is known that cellobiose acts as an inhibitor of cellobiohydrolases and endoglucanases and thereby reduces the rate of hydrolysis for the entire cellulase system. Therefore, the use of increased  $\beta$ -glucosidase activity to quickly convert cellobiose into glucose would greatly enhance the production of ethanol. To illustrate this point, cytolase 123 and the fermentation product produced by the transformants (normalized to cytolase on a total protein basis) according to the present invention under fermentation conditions were compared for their ability to hydrolyze crude paper fractions composed of 50-60% cellulotics from a fiber fraction (RDF) of municipal solid waste (MSW). Such suspensions were in 50 mM sodium acetate buffer, pH 4.8 to 5.0, and equilibrated at 30°C. The flasks were then dosed with 4% Saccharomyces cerevisiae and sampled periodically to 80 hours. The ethanol production yield was then measured. The following Table II illustrates that increased ethanol production is possible using the increased  $\beta$ -glucosidase preparation from the present invention using municipal solid waste preparations as the cellulosic source.

TABLE II

mg protein	gram cellulose	Grams/Liter Ethanol	
		Cytolase 123	High $\beta$ -Glu Prep
10		2.1	5.0
20		5.3	7.2
30		6.9	8.8
40		8.0	9.3
50		8.5	9.3
60		8.5	9.3

[0099] From Table II it can be clearly seen that the enhanced  $\beta$ -glucosidase preparation prepared according to the present invention enhances the production of ethanol compared to a cytolase 123 control, especially at the lower protein concentrations.

[0100] The detergent compositions of this invention may employ, besides the cellulase composition, a surfactant, including anionic, nonionic and amphoteric surfactants, a hydrolase, building agents, bleaching agents, bluing agents and fluorescent dyes, caking inhibitors, solubilizers, cationic surfactants and the like. All of these components are known in the detergent art. For a more thorough discussion, see U.S. Application Serial No. 07/593,919 entitled "Trichoderma reesei Containing Deleted Cellulase Genes and Detergent Compositions Containing Cellulases Derived Therefrom", and U.S. Serial No. 07/770,049, filed October 4, 1991 and entitled "Trichoderma reesei Containing Deleted and/or enriched Cellulase and other enzyme Genes and Cellulase Compositions Derived Therefrom" both of which are incorporated herein by reference in their entirety.

[0101] The detergent compositions contain enhanced levels of  $\beta$ -glucosidase or altered  $\beta$ -glucosidase. In this regard, it really depends upon the type of product one desires to use in detergent compositions to give the appropriate effects.

[0102] Preferably the cellulase compositions are employed from about 0.00005 weight percent to about 5 weight percent relative to the total detergent composition. More preferably, the cellulase compositions are employed from about 0.01 weight percent to about 5 weight percent relative to the total detergent composition and even more preferably, from about 0.05 to about 2 weight percent relative to the total detergent composition.

[0103] Moreover, the present invention also contemplates the use of the  $\beta$ -glucosidase nucleotide sequence of T. reesei to design various probes for the identification of the extracellular  $\beta$ -glucosidase gene in other filamentous fungi. In this regard, the entire nucleotide sequence of the bgl1 gene can be used or a portion thereof to identify and clone out the equivalent genes from other filamentous fungi. The sources of filamentous fungi include those fungi from the genus Trichoderma, Aspergillus, Neurospora, Humicola, Penicillium and the like. More particularly, the preferred species include Trichoderma reesei, Trichoderma viridae, Aspergillus niger, Aspergillus oryzae, Neurospora crassa, Humicola grisea, Humicola insolens, Penicillium pinophilum, Penicillium oxalicum, Aspergillus phoenicis, Trichoderma koningii and the like. Due to the species homology of the bgl1 gene, filamentous fungi equivalent genes are easily

identified and cloned. Indicative of this are Figures 10A and 10B which illustrate autoradiograph of *A. nidulans* and *N. crassa* (Figure 10A) and *H. grisea* (Figure 10B) DNA digested with *Hind* III and *Eco* RI and further were blotted and probed with a  $P^{32}$  labeled *Hind* III 6.0 kb *bgl1* DNA fragment containing the *bgl1* gene of *T. reesei*. These autoradiographs clearly illustrate that a DNA fragment containing the *bgl1* gene of *T. reesei* can be used to identify the extracellular

*bgl1* gene in other fungi.

[0104] Thus the *bgl1* gene of other filamentous fungi may be cloned by the methods outlined above using the  $P^{32}$  labelled *T. reesei* *bgl1* gene as a probe. Once the genes of other filamentous fungi are cloned, they can be used to transform the filamentous fungi from which the gene was derived or other filamentous fungi to overproduce  $\beta$ -glucosidase by the methods described above.

[0105] In order to further illustrate the present invention and advantages thereof, the following specific examples are given, it being understood that the same are intended only as illustrative and in nowise limitative.

#### Example 1

##### Isolation of Total RNA from *Trichoderma reesei*

[0106] A *Trichoderma reesei* culture which over produces cellulases was specifically induced for cellulase using sophorose, a  $\beta$ , 1-2 diglucoside as described by Gritzali, 1977. The starting strain of *Trichoderma reesei* is a cellulase over-production strain (RL-P37) developed by mutagenesis by the methods described by Sheir-Neiss, G. and Montencourt, B.S., *Appl. Microbiol. Biotechnol.*, Vol. 20 (1984) pp. 46-53. A mycelial inoculum of *T. reesei*, from growth on potato dextrose agar (Difco), was added into 50 ml of *Trichoderma* basal medium containing 1.40 grams/liter  $(NH_4)_2SO_4$ , 2.0 grams/liter  $KH_2PO_4$ , 0.30 grams/liter  $MgSO_4$ , 0.30 grams/liter urea, 7.50 grams/liter BactoPeptone, 5.0 ml/liter, 10% Tween - 80, 1.0 ml/liter trace elements-EFG, pH 5.4, which was filtered through a 0.2 micron filter in a 250 ml baffled flask. This culture was incubated at 30°C for 48 hours with vigorous aeration. Five milliliter aliquots were taken from the culture and added to 25 ml of fresh basal medium in seven 250 ml flasks. These were subsequently grown for 24 hours at 30°C. All cultures were centrifuged in a benchtop clinical centrifuge at 2400 x g for 10 minutes. The mycelial pellets were washed three times in 50 ml of 17 mM  $KHPO_4$  buffer (pH 6.0). Lastly, the mycelia were suspended in six flasks containing 50 ml of 17 mM  $KHPO_4$  buffer with the addition of 1 mM sophorose and a control flask containing no sophorose. The flasks were incubated for 18 hours at 30°C prior to harvesting by filtration through Miracloth (Calbiochem). The excess medium was then squeezed out and the mycelial mat was placed directly into liquid nitrogen and may be stored at -70°C for up to one month. The frozen hyphae were then ground in an electric coffee grinder that was prechilled with a few chips of dry ice until a fine powder was obtained. The powder was then added to about 20 ml of an extraction buffer containing 9.6 grams of p-aminosalicylic acid dissolved in 80 ml of DEP-treated water, 1.6 grams of triisopropylphenyl sulfonic acid dissolved in 80 ml of DEP-treated water, 24.2 grams

Tris-HCl, 14.6 grams NaCl, 19.09 grams EDTA, which was diluted to 200 ml total volume with DEP-treated water and the pH was adjusted to 8.5 with NaOH. After addition of the extraction buffer, 0.5 volumes of TE-saturated phenol was also added thereto, and the extraction mixture was placed on ice. One quarter volume of chloroform was then added to the extraction mixture, and the mixture was shaken for two minutes. The phases were then separated by centrifugation at 2500 rpm. The aqueous phase was removed and placed in a centrifuge tube, which contained a few drops of phenol in the bottom of said tube. The tube was placed on ice. The organic phase was then reextracted with 2.0 ml of extraction buffer and placed in a 68°C water bath for 5 minutes to release the RNA trapped in polysomes and at the interface of the extraction mixture. The extracted mixture was then centrifuged, and the aqueous phase removed and pooled with the other aqueous fraction.

[0107] The entire aqueous fractions were then extracted with phenol-chloroform (1:1 v/v) for 4 to 5 times until there was no longer any protein seen visually at the interface. Then 0.1 volume of 3 M sodium acetate, pH 5.2 (made with DEP water and autoclaved) and 2.5 volumes of 95% was added to the organic extracts, and the extracts were frozen at -20°C for 2 to 3 hours. Alternatively, the RNA was precipitated using 2 M lithium acetate. The RNA was then pelleted by centrifugation at 12,000 rpm for 20 minutes. The pelleted RNA was then resuspended in DEP-water with an RNase inhibitor to a final concentration of 1 unit per  $\mu$ l. To determine whether the genes encoding the enzymes were being induced, total RNA was analyzed.

##### Analysis of Total RNA Preparation

[0108] To confirm whether the genes encoding the enzymes of the cellulase complex were being induced, total RNA was analyzed by Northern blotting as described by Sambrook et al, *supra* using a  $P^{32}$  fragment of the *T. reesei* *cbh2* gene as a probe. The *cbh2* clone was isolated using the methods described by Chen et al in "Nucleotide Sequence and Deduced Primary Structure of Cellobio-hydrolase II from *Trichoderma reesei*", *Biotechnology*, Vol. 5 (March 1987), incorporated herein by reference. Site directed mutagenesis (Sambrook et al., *supra*) was performed on the *cbh2* clone



and a Bgl II site was placed at the exact 5' end of the opening reading frame and an Nhe I site at the exact 3' end. The Bgl II/Nhe I coding sequence was then cloned into a pUC218 phagemid. For use as a probe, the cbh2 fragment was digested with Bgl II/Nhe I and isolated by gel electrophoresis. The results indicated that the level of cbh2 specific mRNA reached a peak at 14-18 hours post induction. The total RNA from 14, 18 and 22 hours was then pooled.

## Example 2

### Purification of Polyadenylated mRNA

[0109] mRNA was then isolated from the pooled fraction of total RNA set forth above using oligo (dT) cellulose chromatography. Oligo(dT) cellulose (type 3 from Collaborative Research, Lexington, MA) is first equilibrated with Oligo (dT) binding buffer containing 0.01 M Tris-HCl, pH 7.5, 0.5 M NaCl, and 1 mM EDTA, then aliquots of 25-300 mg were added to 1.5 ml microfuge tubes. RNA dissolved in 1 ml of binding buffer was added and allowed to bind for 15 min. with gentle shaking. The suspensions were centrifuged at 1500 g for 3-5 min., washed 3-5 times with 1 ml of binding buffer, and then washed 3 times with 400 µl of elution buffer containing 0.01 M Tris-HCl, pH 7.5, and 1 mM EDTA. The eluates were pooled, readjusted to 0.5 M NaCl, rebound, and reeluted with three washes of elution buffer. The final three elution buffer washes were pooled and mRNA was recovered by ethanol precipitation.

### Analysis of Total RNA and polyadenylated mRNA

[0110] Total RNA and the polyadenylated RNA were fractionated on 1% formaldehyde-agarose gels using 10 µg of RNA for each lane, blotted to Nytran<sup>R</sup> membranes and analyzed by the Northern blot method described by Thomas in "Hybridization of denatured RNA and Small DNA fragments transferred to Nitrocellulose", Proc. Natl. Acad. Sci. USA, Vol. 77 (1980), pp. 5201-5205.

[0111] Briefly, this procedure involves denaturing RNA (up to 10 µg/8 µl reaction) by incubation in 1 M glyoxal/50% (vol/vol) Me<sub>2</sub>SO/10 mM sodium phosphate buffer, pH 7.0 at 50°C for 1 hr. The reaction mixture was cooled on ice and 2 µl of sample buffer containing 50% (vol/vol) glycerol, 10 mM sodium phosphate buffer at 7.0 and bromophenol blue was added. The samples were electro-phoresed on horizontal 1% formaldehyde-agarose gels in 10 mM phosphate buffer, pH 7.0 at 90 v for 6 hours.

[0112] The glyoxylated RNA was transferred from agarose gels to nitrocellulose by using 3 M NaCl/0.3 M trisodium citrate (20X NaCl/cit). After electrophoresis, the gel was placed over two sheets of Whatman 3 MM paper which was saturated with 20X NaCl/cit. Nitran<sup>R</sup> membrane was wetted with water, equilibrated with 20X NaCl/cit and laid over the gel. The gel was then covered with two sheets of Whatman 3 MM paper and a 5 to 7 cm layer of paper towels, a glass plate and a weight. Transfer of the RNA was completed in 12-15 hours. The blots were then dried under a lamp and baked in a vacuum for over 2 hrs. at 80°C.

[0113] The membranes were probed with a cbh2 probe to verify that the polyadenylated mRNA pool contained cbh2 mRNA and by inference the genes encoding the enzymes of the cellulase complex were indeed induced.

## Example 3

### Synthesis of cDNA

#### A. First Strand Synthesis

[0114] Synthesis of cDNA was performed using the BRL cDNA Synthesis System<sup>R</sup> (Bethesda Research Laboratories, Md.) according to the instructions of the manufacturer. To a sterile, DEPC-treated tube in ice was added 10 µl of 5X First Strand Buffer containing 250 mM Tris-HCl, pH 8.3, 375 mM KCl, 15 mM MgCl<sub>2</sub>, 50 mM DTT, 2.5 µl 10 mM dNTP Mix (10 mM dATP, 10 mM dCTP, 10 mM dGTP, 10 mM dTTP), 5 µl Oligo (dT)<sub>12-18</sub> (0.5) mg/ml, 10 µl of mRNA at 0.5 mg/ml and 20 µl diethylpyrocarbamate(DEPC)-treated water to create a final composition containing 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl<sub>2</sub>, 10 mM dithiothreitol, 500 µM each dATP, dCTP, dGTP and dTTP, 50 µg/ml Oligo (dT)<sub>12-18</sub>, 100 µg/ml polyadenylated RNA and 10,000 U/ml cloned M-MLV reverse transcriptase. A control run was also run simultaneously using 10 µl of a 2.3 kb control RNA (0.5 mg/ml) in lieu of the mRNA.

[0115] The reaction was initiated by adding 2.5 µl of Molony murine leukemia virus (M-MLV) reverse transcriptase (100 Units/µl) to the mRNA tube and the control RNA. The samples were mixed. All reaction tubes were incubated at 37°C for one hour and then placed on ice.

[0116] A small aliquot from the reaction mixture was run on a gel to confirm its presence and quantity. The yield obtained was about 2-6 µg.



## B. Second Strand Synthesis

[0117] To the control tube on ice after first strand synthesis was added 230.6 µl DEPC-treated water, 6 µl 10 mM dNTP mix, 32 µl 10X second strand buffer containing 188 mM Tris-HCl, pH 8.3, 906 mM KCl, 100 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 46 mM MgCl<sub>2</sub>, 37.5 mM dithiothreitol, 1.5 mM NAD, 8 µl *E. coli* DNA Polymerase I (10 µl/µl), 1.4 µl *E. coli* RNase H and 1 µl *E. coli* DNA ligase (100 units).

[0118] To the first strand synthesis of the sample was added on ice 289.5 µl of DEPC-treated water, 7.5 µl 10 mM dNTP mix, 40 µl 10X second strand buffer, 10 µl *E. coli* DNA Polymerase I, 1.75 µl *E. coli* RNaseH and 1.25 µl *E. coli* DNA ligase, to create a final composition containing 25 mM Tris-HCl (pH 8.3), 100 mM KCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 5 mM MgCl<sub>2</sub>, 250 µM each dATP, dCTP, dGTP, dTTP, 0.15 mM NAD, 5 mM Dithiothreitol, 250 U/ml DNA Polymerase I, 8.5 U/ml RNase H, and 30 U/ml DNA Ligase. Both the control tube and the sample tube were vortexed gently and incubated for 2 hours at 16°C. After incubation, both tubes were placed on ice.

[0119] The sample tube was then extracted with 415 µl of phenol and ethanol precipitated. The pellet was dissolved in 200 µl of sterile TE buffer (10 mM Tris-HCl pH 7.5, 1 mM Na<sub>2</sub>EDTA) and reprecipitated from 7.5 M ammonium acetate with ethanol.

[0120] An aliquot of the sample was further analyzed by gel electrophoresis to check for purity. The yield of the synthesis was about 4.0 µg.

[0121] The remaining control sample was further extracted with phenol and ethanol precipitated as described above for the sample. After dissolving the pellet in 200 µl of sterile TE buffer, reprecipitating the sample from ammonium acetate with ethanol, and redissolving the dry pellet in 20 µl of sterile TE buffer, 2 µl of the solution was then further analyzed by gel electrophoresis to check for purity.

### Example 4

#### Amplification of bgl1 cDNA Sequences

[0122] Amplification of the cDNA fragments encoding a portion of the *T. reesei* β-glucosidase gene, *bgl1*, was performed using the polymerase chain reaction (PCR) method with Taq<sup>R</sup> polymerase and a Perkin Elmer Cetus Thermal Cycler<sup>R</sup>.

[0123] The reaction mixture was formed by mixing 76 µl deionized water, 10 µl of a 10X mixture of buffer containing 166 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 670 mM Tris-HCl, pH 8.8, 67 mM MgCl<sub>2</sub>, 67 µM EDTA, 10 mM β-mercaptoethanol, 10 µl dimethylsulfoxide and 1.7 mg/ml BSA diluted to a total volume of 1.0 ml with deionized water, 8 µl of 2 dNTPs (each), 1 µl 5' oligonucleotide primer, 1 µl 3' internal oligonucleotide primer, 1.0 µg cDNA diluted in 3 µl deionized water, and 1 µg Taq<sup>R</sup> polymerase.

[0124] The amplification method consists of an initial denaturing cycle at 95°C for 10 minutes, followed by a two minute annealing step at 50°C and a 10 minute polymerization cycle at 65°C, for an additional 30 cycles.

#### A. Oligonucleotide Primers

[0125] The oligonucleotide primers used to amplify the cDNA fragment encoding the *T. reesei* *bgl1* gene were designed based on the degeneracy of the genetic code for the selected amino acids for an N terminal region of the *bgl1* gene and an internal oligonucleotide. The 5' oligonucleotide primer consisted of the sequence:

5' GCI GTI CCT CCT GCI GG 3'

wherein I = inosine.

[0126] The internal 3' oligonucleotide primer consisted of a pool of 16 x 21 oligonucleotides. This pool was based on various derivations of the following sequences:

5' GTT G/ATT ICC G/ATT G/AAA G/ATC TGT 3'

### Example 5

#### Subcloning of PCR Generated Fragments

[0127] Ninety µl of each reaction mix was fractionated on 4% polyacrylamide gels in 1X TBE, the major band was

excised and eluted from the gel slice as described by Sambrook et al, supra. The eluted DNA fragment was precipitated in ethanol and resuspended in 15 µl of TE buffer (10 mM Tris, 1 mM EDTA). Each 1-2 µg DNA fragment was then treated with 0.5 mM ATP and T<sub>4</sub> polynucleotide kinase to phosphorylate the 5' end of each fragment following by the procedures of Sambrook et al, supra. Blunt ends were generated by adding 3 µl of 10X T<sub>4</sub> polymerase buffer (330 mM Tris-acetate at pH 7.9, 660 mM potassium acetate, 100 mM magnesium acetate, 1 µl of 2.5 mM dNTPS, 1 µl of T<sub>4</sub> DNA polymerase and 5 µl of distilled water). The blunt-end reaction mixture was then incubated at 37° for 60 minutes. The reaction was stopped by addition of EDTA to a final concentration of 1 mM EDTA and the sample was further heated for 10 minutes at 65°C.

[0128] The blunt-end DNA fragments were then ligated with SmaI cleaved and dephosphorylated pUC218 which had been infected with M13X07 as described by Sambrook et al, supra. The cloning vectors pUC218 and pUC219 were derived from pUC118 and pUC119 by insertion of the Bgl II, Cla I and Xho I polylinker as described by Korman et al in "Cloning, Characterization, and expression of two α-amylase genes from Aspergillus niger var. awamori", Current Genetics, Vol. 17, pp. 203-212, (1990).

[0129] The aforescribed phagemid was then used to transform E. coli strain JM101 as described by Yarnisch et al in "Improved M13 phage cloning vectors and host strains: nucleotide sequence of the M13mp18 and pUC19 Vectors", Gene, Vol. 1197, pp. 103-119 (1985).

#### Example 6

##### Isolation of cDNA Subcloned Fragment

[0130] The transformed strain was inoculated in 1.5 ml of 2YT broth in a tube which had been previously inoculated with 15 µl of saturated E. coli JM101. The culture was grown for 8 hours under shaking at 37°C.

[0131] The culture mixture was then spun at 6000 rpm for 5 minutes, and the supernatant was poured off into another tube. To the supernatant 300 µl of 2.5 M NaCl, 20% PEG was added, and the solution was mixed. The mixture was then incubated at room temperature for 15 minutes.

[0132] The solution was then spun for 5 minutes in a microfuge, and the supernatant was aspirated off. The solution was vortexed once again, and the supernatant was further aspirated off.

[0133] 100 µl of equilibrated phenol was added to the tube, and the tube was vortexed. 100 µl of chloroform was added, and once again the tube was vortexed. The tube was heated at 55°C for 5 minutes, mixed, and microfuged an additional 5 minutes.

[0134] 160 µl of the supernatant was then pipetted off and transferred to a clear tube. 20 µl of 1N NaOAC, pH 4.5, and 400 µl of 95% ETOH were added to the supernatant, and the solution was mixed and frozen on dry ice for 5 minutes. The tube was then spun for an additional 15 minutes, and the supernatant was aspirated off.

[0135] 1000 µl of 70% ethanol was added to the tube, and the tube was spun for an additional 2 minutes and reaspirated. The mixture was spun once more under vacuum for 4 minutes, and the pellet was resuspended in 15 µl TE buffer.

#### Example 7

##### Determination of the Nucleotide Sequence of 700 bp cDNA fragment

[0136] The nucleotide sequence of the 700 bp cDNA fragment was determined using the dideoxy DNA sequencing method described by Sanger et al, "DNA Sequencing with chain terminating inhibitors", Proc. Natl. Acad. Sci. U.S.A., Vol. 74 (1977), p. 5463, using the Sequenase<sup>R</sup> reagent kit (U.S. Biochemicals).

#### Example 8

##### Analysis of bgl1 gene

##### A. Sequence Analysis

[0137] Nucleotide sequencing was done by the dideoxy chain termination method of Sanger et al (1977) using the Sequenase<sup>R</sup> reagent kit (U.S. Biochemicals).

##### B. Amino Acid Sequencing

[0138] A 2.5-nmol sample of the reduced and carboxymethylated β-glucosidase preparation purified (per Chirico and

Brown, European Journal of Biochem., Vol. 165, pp. 333 et seq.) was subjected to N-terminal sequencing on a proprietary multiphase sequencer.

[0139] To a sample of  $\beta$ -glucosidase, Endo-Lys C protease was added to 1% of the total protein and the mixture incubated for 1 hour at 37°C or the protein sample was subject to cyanogen bromide treatment. An equal volume of HPLC solution A (0.05% TEA/0.05% TFA in water) was added to stop the reaction. The resulting CNBr and Endo-Lys C fragments were separated by chromatography on a Brownlee C-4 column using a linear gradient of 0-100% HPLC solution B (0.05% TEA/0.05% TFA in n-propanol) at a rate of 1% per minute. Several peaks were collected for amino acid sequencing and the data are denoted in Fig. 1.

## 10 Example 9

### Identification of *bgl1* gene from *T. reesei*

[0140] The 700 bp *bgl1* cDNA fragment was then labelled with  $^{32}\text{P}$  using methods described by Sambrook et al, *supra*.

15 [0141] Genomic DNA from *T. reesei* was prepared by filtering a 24-36 hour culture of *T. reesei* through Miracloth and freezing the mycelia obtained from the culture medium. The frozen mycelia were then ground into a fine powder and 22 mls of TE, and 4 mls of 20% SDS were added to the powdered mycelia and mixed. 10 ml of phenol and chloroform was added to the mixture prior to centrifugation and removal of the aqueous phase. 200  $\mu\text{l}$  of 5 mg/ml proteinase K was added to the organic extract, and the mixture was incubated for 20 minutes at 55°C. The DNA was then further  
20 extracted by methods known in the art using chloroform/phenol extraction followed by ethanol precipitation. The isolated DNA was then treated with 1  $\mu\text{g}$  of heated ribonuclease A (100°C for 15 minutes) per 20  $\mu\text{g}$  of genomic DNA in TE buffer at 37°C for 30 minutes, then cooled to room temperature. The genomic DNA from *T. reesei* was then cut singly or in combination with a variety of restriction enzymes such as *Eco* RI, *Hind* III and the like, Southern blotted and hybridized with the  $^{32}\text{P}$  labelled 700 bp cDNA fragment of the *bgl1* gene as a probe. From this analysis it was determined  
25 that *Hind* III was the restriction enzyme of choice used to locate the  $\beta$ -glucosidase gene.

[0142] 10 to 20 units of *Hind* III per milligram of genomic DNA was added to the DNA and then the DNA was extracted with phenol-chloroform to remove protein. The treated DNA was then alcohol precipitation and resuspended to 2 grams/liter in TE buffer.

30 [0143] 4  $\mu\text{l}$  samples from the *Hind* III digestion of genomic DNA were loaded on a 1% agarose gel and fractionated electrophoretically. The gel was then Southern blotted and probed with the  $^{32}\text{P}$  700 bp cDNA probe. A 6.0 kb band was identified on the Southern blot of *Hind* III digested genomic DNA from *T. reesei*.

[0144] The remaining *Hind* III genomic DNA was then subjected to a preparative gel electrophoresis and fragments ranging from 5 kb to 7 kb were then electroeluted from the agarose gel and cloned into *Hind* III digested pUC218. The resulting plasmids were used to transform *E. coli* JM101 to create a library. Then the library was screened by colony  
35 hybridization using  $^{32}\text{P}$  labelled 700 bp *bgl1* cDNA as a probe to identify those colonies containing DNA coding for the *bgl1* gene.

[0145] The positive colonies from the transformation were then picked and the DNA isolated therefrom by phenol: chloroform extraction and ethanol precipitation, described by Sambrook et al, *supra*.

40 [0146] The isolated DNA from the positive colonies was digested both singly and in various combinations with the following restriction enzymes: *Hind* III, *Eco* RI, *Sst* I, *Kpn* I, *Bam* HI, *Xho* I, *Bgl* II, *Cla* I, *Xba* I, *Sal* I, *Pst* I, *Sph* I, *Bal* I, and *Pvu* II. The digestions were subjected to agarose gel electrophoresis, and the resultant banding pattern was used to construct a restriction map of the cloned 6.0 kb genomic DNA. The same agarose gel was Southern blotted and probed with the  $^{32}\text{P}$  labelled 700 bp *bgl1* cDNA to identify which genomic restriction fragments shared homology with the *bgl1* cDNA. The mapping experiments confirmed that the entire *bgl1* gene is contained on the genomic *Hind*  
45 III clone. *Pvu* II and *Bal* I restriction fragments which ranged in size from 600 bp to 1500 bp hybridized with the 700 bp DNA *bgl1* clone and were chosen for subcloning into pUC218 phagemid. After cloning these fragments into the phagemid, the *Pvu* II and *Bal* I subclones were then sequenced using the dideoxy chain termination method of Sanger et al (1977). It was then determined from this sequencing that the overlapping sequences of the subclones aligned with a single contiguous sequence totaling 3033 bp within which the nucleotide sequence was determined on both  
50 strands.

## Example 10

### Construction of pSAS $\beta$ -glu

55 [0147] The starting vector for the construction of pSAS $\beta$ -glu was the plasmid pSAS. pSAS was constructed in the following way. pUC100 (a commercially available plasmid vector) was digested with the restriction enzyme *Sma*I and the 5' phosphate groups subsequently removed by digestion with calf intestinal alkaline phosphatase. The linear v ctor

fragment was purified from undigested vector and protein by agarose gel electrophoresis followed by isolation of the linear vector DNA from the isolated gel slice by electroelution. The amdS gene was isolated as a 2.4 kb SstI restriction fragment following separation from the vector sequences (contained in - Hynes, M.J., Corrick, C.M., and King, J.A., "Isolation of genomic clones containing the amdS gene of *Aspergillus nidulans* and their use in the analysis of structural and regulatory mutations", *Mol. Cell. Biol.*, Vol. 3 (1983), pp. 1430-1439). The 2.4 kb SstI amdS fragment and the 2.7 kb pUC100 vector fragment were then ligated together (Sambrook et al., *supra*) and the ligation mix transformed and propagated in the *E. coli* host strain, JM101.

[0148] pSAS $\beta$ -glu was constructed by digesting pSAS with the restriction enzyme Hind III, and purifying the linear fragment as described above. Into this Hind III treated pSAS vector fragment was ligated a 6.0 kb Hind III fragment of *T. reesei* genomic DNA that contained all of the coding region of the bgl1 gene along with sequences necessary for the genes transcription and translation.

#### Example 11

##### Preparation of BGL1 Deletion Vector

[0149] The gene replacement vector pUC $\Delta\beta$ -Glu A/R pyr, illustrated in Figure 3B, was constructed by cloning a 6.0 kb genomic HindIII fragment, known to contain the entire bgl1 gene, into the polylinker of pUC218 which had been cut with HindIII and the ends dephosphorylated with calf intestinal alkaline phosphatase. The coding region for the bgl1 gene was then removed from this plasmid by digesting the plasmid with Apal and EcoRV at unique Apal and EcoRV restriction sites situated at the very 5' and 3' end of the bgl1 open reading frame and isolating the linear plasmid DNA. The restriction site ends were made blunt with T4 DNA polymerase. This plasmid was then ligated with an isolated 2412 bp Hind III/Bam HI restriction fragment containing the pyrG gene from *Aspergillus niger* (Hartingsreldt et al., *Mol. Gen. Genet.* 206:71-75 (1987) in which the restriction ends were made blunt by treatment with T4 DNA polymerase to create pUC $\Delta\beta$ Glu A/R pyr (Fig. 3B).

#### Example 12

##### Isolation of Protoplasts

[0150] Mycelium was obtained by inoculating 100 ml of YEG (0.5% yeast extract, 2% glucose) in a 500 ml flask with about  $5 \times 10^7$  *T. reesei* cells. The flask was then incubated at 37°C with shaking for about 16 hours. The mycelium was harvested by centrifugation at 2,750 x g. The harvested mycelium were further washed in 1.2 M sorbitol solution and resuspended in 40 ml of Novozym<sup>R</sup>, which is the tradename for a multi-component enzyme system containing 1,3-alpha-glucanase, 1,3-beta-glucanase, laminarinase, xylanase, chitinase and protease from Novo Biolabs, Danbury, Ct., solution containing 5 mg/ml Novozym<sup>R</sup> 234; 5 mg/ml MgSO<sub>4</sub>·7H<sub>2</sub>O; 0.5 mg/ml bovine serum albumin; 1.2 M sorbitol. The protoplasts were removed from cellular debris by filtration through Miracloth (Calbiochem Corp.) and collected by centrifugation at 2,000 x g. The protoplasts were washed three times in 1.2 M sorbitol and once in 1.2 M sorbitol, 50 mM CaCl<sub>2</sub>, centrifuged and resuspended. The protoplasts were finally resuspended at a density of  $2 \times 10^8$  protoplasts per ml of 1.2 M sorbitol, 50 mM CaCl<sub>2</sub>.

#### Example 13

##### Transformation of Fungal Protoplasts with pSAS $\beta$ -glu

[0151] 200  $\mu$ l of the protoplast suspension prepared in Example 12 was added to 20  $\mu$ l (20  $\mu$ g) of pSAS $\beta$ -glu in TE buffer (10 mM Tris, pH 7.4; 1 mM EDTA) and 50  $\mu$ l of a polyethylene glycol (PEG) solution containing 25% PEG 4000, 0.6 M KCl and 50 mM CaCl<sub>2</sub>. This mixture was incubated on ice for 20 minutes. After this incubation period 2.0 ml of the above-identified PEG solution was added thereto, the solution was further mixed and incubated at room temperature for 5 minutes. After this second incubation, 4.0 ml of a solution containing 1.2 M sorbitol and 50 mM CaCl<sub>2</sub> was added thereto and this solution was further mixed. The protoplast solution was then immediately added to molten aliquot's of Vogel's Medium N (3 grams sodium citrate, 5 grams KH<sub>2</sub>PO<sub>4</sub>, 2 grams NH<sub>4</sub>NO<sub>3</sub>, 0.2 grams MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.1 gram CaCl<sub>2</sub>·2H<sub>2</sub>O, 5  $\mu$ g  $\alpha$ -biotin, 5 mg citric acid, 5 mg ZnSO<sub>4</sub>·7H<sub>2</sub>O, 1 mg Fe(NH<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O, 0.25 mg CuSO<sub>4</sub>·5H<sub>2</sub>O, 50  $\mu$ g MnSO<sub>4</sub>·4H<sub>2</sub>O containing an additional 1% glucose, 1.2 M sorbitol and 1% agarose. The protoplast/medium mixture was then poured onto a solid medium containing the same Vogel's medium as stated above containing in addition acetamide as a nitrogen source. Since *T. reesei* does not contain a functional equivalent to the amdS gene only transformants will grow on this medium. These colonies were subsequently transferred and purified on a solid Vogel's medium N containing as an additive, 1% glucose. The bgl1 gene inserted transformant strain is called A83pSAS $\beta$ Glu.

[0152] Stable transformants can be distinguished from unstable transformants by their faster growth rate and the formation of circular colonies with a smooth rather than ragged outline on solid culture medium. Additionally, in some cases, a further test of stability can be made by growing the transformants on solid non-selective medium, harvesting the spores from this culture medium and determining the percentage of these spores which will subsequently germinate and grow on selective medium.

[0153] Figure 6 is an autoradiograph of a Southern blot using the  $P^{32}$  labelled 700 bp fragment as a probe, of the different transformants with enhanced copies of the *bgl1* gene (lanes 1-8) using genomic *T. reesei* from an overproducing strain digested with *Hind* III as a control (lane 9). This autoradiograph clearly shows that the transformants contained enhanced amount of the *bgl1* gene compared with the control.

[0154] Figure 4 is an autoradiograph of a Northern blot of RNA isolated from one of the transformed strains (lane A) produced by the present invention following induction with sorbose illustrating a corresponding increase in the levels of *bgl1* message when compared to the parental strain of *T. reesei* (lane B).

[0155] Besides visual analysis of the transformants, quantitative analysis was also completed by cutting the appropriate bands out of the Nytran<sup>R</sup> membrane and counting the radioactive label present therein in a scintillation counter. This experiment was performed to obtain a more precise estimate of the relative amounts of message as shown in Table III below:

TABLE III

CPM	Parental <i>Trichoderma reesei</i> strain	Transformed <i>Trichoderma reesei</i> strain
CPM $\beta$ -glu message	14.4	25.4
CPM CBHII	227.1	95.2
CPM $\beta$ -glu/CBHII	0.0634	0.2668

[0156] Table III illustrates that the transformant produced by the process of the present invention has extra  $\beta$ -glucosidase mRNA and hence an increase in  $\beta$ -glucosidase enzyme resulting in an increase in specific activity.

#### Example 14

##### Transformation of Fungal Protoplasts with pUC $\Delta\beta$ Glu A/R *pyr4*

[0157] Mutants of *T. reesei* lacking the coding sequence for the extracellular  $\beta$ -glucosidase gene, *bglI*, were obtained by a targeted gene replacement event. pUC $\Delta\beta$ Glu A/R *pyr4* plasmid was digested with *Hind* III to obtain a linear *Hind* III fragment in which the *bglI* coding sequences were replaced with the *pyrG* gene from *Aspergillus niger*. Protoplasts were transformed with the linear DNA fragment containing the *bgl1* flanking sequences and the *pyr4* by the methods of Examples 12 and 13. The deletion transformants were called  $\Delta 12$  and  $\Delta 36$ . After transformation, the protoplast solution was then added to molten aliquots of Vogel's Medium N containing an additional 1% glucose, 1.2 M sorbitol and 1% agarose. The protoplast/medium mixture was then poured into a solid medium containing the same Vogel's medium N. No uridine was present in the medium and therefore only transformed colonies were able to grow as a result of complementation of the *pyr4* mutation of the *T. reesei* strain RL-P37 by the wild type *pyr4* gene inserted in the DNA fragment. Stable transformants were then selected by the method recited in Example 13.

#### Example 15

##### Analysis of the Transformants

[0158] The transformants were analyzed for the presence or absence of the *bgl1* gene using the 700 bp cDNA probe recited above. The transformants were digested using *Hind* III. Total genomic DNA from selected transformants was digested with *Hind* III restriction enzyme, run on a 1% agarose gel, transferred to Nitran<sup>R</sup> membrane and probed with a  $P^{32}$  labelled 700 bp cDNA recited above and visualized by autoradiography on X-ray film. The results of this analysis are set forth in Figure 5A illustrate that the transformants ( $\Delta 12$  and  $\Delta 36$ ) did not contain a band corresponding to the *bgl1* gene whereas the wild type strain (RL-P37, i.e., P-37) did.

[0159] mRNA isolated from the transformants of Example 14 and analyzed on a Northern blot, as in Example 2. As indicated in Figure 5B, Northern blot analysis using the  $P^{32}$  labelled 2.2 Kb *Apal/EcoRV* *bgl1* probe indicated that *bgl1* specific mRNA was present in *T. reesei* RL-P37 *pyrG69* and is absent in the transformants  $\Delta 12$  and  $\Delta 36$ .

[0160] Protein was recovered as per Example 8 above and then analyzed for the presence of  $\beta$ -glucosidase by use of polyclonal antibodies (from rabbits challenged with pure  $\beta$ -glucosidase) tagged with horseradish peroxidase to permit

detection. The antibodies were used to identify pure  $\beta$ -glucosidase (100 ng -- Column A; 1000 ng -- Column B); cellulase produced from wild type *T. reesei* (Column C); and from cellulase produced by a *T. reesei* strain genetically engineered to delete the  $\beta$ -glucosidase gene (Column D). The results of this analysis are set forth in Figure 5C and show that only Column D did not contain  $\beta$ -glucosidase.

[0161] While the invention has been described in terms of various preferred embodiments, the skilled artisan will appreciate that various modifications, substitutions, omissions, and changes may be made without departing from the scope thereof. Accordingly, it is intended that the scope of the present invention be limited solely by the scope of the following claims, including equivalents thereof.

## Claims

1. A process for modifying the expression of extracellular  $\beta$ -glucosidase in a filamentous fungus comprising transforming said fungus with an expression vector containing a fungal DNA sequence which:
  - (a) is capable of enhancing the expression of extracellular  $\beta$ -glucosidase through the presence of at least one additional copy of a fungal  $\beta$ -glucosidase gene; or,
  - (b) encodes an altered extracellular  $\beta$ -glucosidase i.e. an enzyme having an amino acid sequence which has been altered with respect to that encoded by the bgl1 gene derived from *Trichoderma reesei* by manipulating said bgl1 DNA sequence.
2. A process for enhancing the expression of extracellular  $\beta$ -glucosidase according to claim 1(a), wherein said expression vector comprises all of a coding region of a fungal  $\beta$ -glucosidase gene and sequences necessary for the  $\beta$ -glucosidase gene's transcription and translation.
3. The process according to claim 1 or claim 2, wherein said filamentous fungus is selected from the genus of *Trichoderma*, *Aspergillus*, *Neurospora*, *Humicola* and *Penicillium*.
4. The process according to claim 3, wherein said filamentous fungus is *Trichoderma reesei*, *Trichoderma viridae*, *Trichoderma koningii*, *Aspergillus niger*, *Aspergillus nidulans*, *Aspergillus wentii*, *Aspergillus oryzae*, *Aspergillus phoenicis*, *Neurospora crassa*, *Humicola grisea*, *Penicillium pinophilum* or *Penicillium oxalicum*.
5. The process according to any one of the preceding claims, wherein said  $\beta$ -glucosidase gene is a bgl1 gene derived from *Trichoderma reesei*.
6. The process according to claim 5, wherein said bgl1 gene comprises the amino acids encoded by the nucleotide sequence from 311 to 2679 of Figure 1.
7. The process according to any one of the preceding claims, further comprising the step of isolating transformants having modified  $\beta$ -glucosidase expression.
8. The process according to claim 7, further comprising the steps of:
  - (a) culturing said transformants under conditions to permit growth of said transformants; and
  - (b) isolating a recombinant fungal cellulase composition produced from said transformants.
9. The process according to claim 8, wherein said recombinant fungal cellulase composition is isolated by:
  - (a) centrifuging said culture medium containing said transformants to form a supernatant and a pellet; and
  - (b) filtering said supernatant to obtain a recombinant fungal cellulase composition.
10. The process according to claim 9, wherein an antimicrobial agent is added to said recombinant fungal cellulase composition after filtration.
11. The process according to any one of claims 8 to 10, further comprising the step of purifying an expression product from said isolated recombinant fungal cellulase composition.
12. A fungal cellulase composition derived from a filamentous fungus comprising a recombinant altered extracellular

$\beta$ -glucosidase produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(b).

- 5 13. A method of producing glucose from cellulosics or hetero-glycans, comprising use of a recombinant fungal cellulase composition having enhanced  $\beta$ -glucosidase activity produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(a).
- 10 14. A method of degrading the cellulosics present in feedstock, biomass or sludge comprising use of a recombinant fungal cellulase composition having enhanced  $\beta$ -glucosidase activity produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(a).
- 15 15. A method for degrading cellulose to glucose comprising use of a recombinant fungal cellulase composition having enhanced  $\beta$ -glucosidase activity produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(a).
- 20 16. A method of using the recombinant fungal cellulase composition having enhanced  $\beta$ -glucosidase activity produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(a), or purified  $\beta$ -glucosidase produced from the process according to claim 12 in food additives to enhance the flavour and aroma.
- 25 17. A detergent composition comprising a cleaning effective amount of a surfactant and at least 0.0001 weight percent of a recombinant fungal cellulase composition having enhanced  $\beta$ -glucosidase activity produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(a).
- 30 18. A method of using the cellulase composition having enhanced  $\beta$ -glucosidase activity produced by the process of any one of claims 8 to 10 only to the extent that they are dependent on claim 1(a), wherein said method comprises adding said recombinant fungal cellulase composition to yeast and biomass to form a mixture and culturing said mixture at a sufficient temperature and for a sufficient period of time to produce ethanol.
- 35 19. Transformants produced by the process according to claim 7 only to the extent to which it is dependent on claim 1(a).
- 40 20. A method of using transformants having enhanced extracellular  $\beta$ -glucosidase activity of claim 19 only to the extent to which it is dependent on claim 1(a), said method comprising adding said transformants to yeast and biomass to form a mixture and culturing said mixture at a sufficient temperature and for a sufficient period of time to produce ethanol.
- 45 21. The method according to claim 18 or claim 20, wherein said yeast is B. clausenii, S. cerevisiae, Cellulolyticus acidothermophilum, C. brassicae, C. lustinaniae, S. uvarum or Schizosaccharomyces pombe.
- 50 22. A nucleotide sequence of a bgl1 gene which entire sequence or a portion is or is not labelled for use as a probe wherein the bgl1 gene has the nucleotide sequence of Figure 1.
23. A method of using a probe comprising a nucleotide sequence according to claim 22, for identifying a  $\beta$ -glucosidase gene from a filamentous fungus.
24. The method according to claim 23 wherein said probe is derived from Trichoderma reesei.
25. The method according to claim 23 or claim 24 wherein one or more probes can be used to identify said  $\beta$ -glucosidase gene from a filamentous fungus.

#### Patentansprüche

1. Verfahren zur Modifizierung der Expression von extrazellulärer  $\beta$ -Glucosidase in einem filamentösen Pilz, umfassend das Transformieren des Pilzes mit einem Expressionsvektor, enthaltend eine Pilz-DNA-Sequenz, welche:
  - (a) fähig zur Erhöhung der Expression von extrazellulärer  $\beta$ -Glucosidase durch die Gegenwart mindestens einer zusätzlichen Kopie eines Pilz- $\beta$ -Glucosidase-Gens ist; oder

(b) eine veränderte extrazelluläre  $\beta$ -Glucosidase codiert, d.h. ein Enzym mit einer Aminosäuresequenz, welche in Bezug auf diejenige, codiert vom bgl1-Gen, abgeleitet aus Trichoderma reesei, durch Manipulieren der bgl1-DNA-Sequenz verändert wurde

- 5 2. Verfahren zur Erhöhung der Expression von extrazellulärer  $\beta$ -Glucosidase gemäß Anspruch 1(a), wobei der Expressionsvektor die gesamte codierende Region eines Pilz- $\beta$ -Glucosidase-Gens und für die Transkription und Translation des  $\beta$ -Glucosidase-Gens notwendige Sequenzen umfasst.
- 10 3. Verfahren nach Anspruch 1 oder Anspruch 2, wobei der filamentöse Pilz gewählt ist aus der Gattung Trichoderma, Aspergillus, Neurospora, Humicola und Penicillium.
- 15 4. Verfahren nach Anspruch 3, wobei es sich bei dem filamentösen Pilz handelt um Trichoderma reesei, Trichoderma viridae, Trichoderma koningii, Aspergillus niger, Aspergillus nidulans, Aspergillus wentii, Aspergillus oryzae, Aspergillus phoenicis Neurospora crassa, Humicola grisea, Penicillium pinophilum oder Penicillium oxalicum.
- 20 5. Verfahren gemäß mindestens einem der vorstehenden Ansprüche, wobei das  $\beta$ -Glucosidase-Gen ein bgl1-Gen, abgeleitet aus Trichoderma reesei, ist.
- 25 6. Verfahren gemäß Anspruch 5, wobei das bgl1-Gen die von der Nukleotidsequenz von Figur 1 von 311 bis 2679 codierten Aminosäuren umfasst.
7. Verfahren gemäß mindestens einem der vorstehenden Ansprüche, ferner umfassend den Schritt des Isolierens von Transformanten mit modifizierter  $\beta$ -Glucosidase-Expression.
- 30 8. Verfahren gemäß Anspruch 7, ferner umfassend die Schritte:
  - (a) Kultivieren der Transformanten unter Bedingungen, um das Wachstum der Transformanten zu gestatten; und
  - (b) Isolieren einer rekombinanten Pilz-Cellulase-Zusammensetzung, hergestellt aus den Transformanten.
- 35 9. Verfahren gemäß Anspruch 8, wobei die rekombinante Pilz-Cellulase-Zusammensetzung isoliert wird durch:
  - (a) Zentrifugieren des Kulturmediums, enthaltend die Transformanten, um einen Überstand und ein Pellet zu bilden; und
  - (b) Filtrieren des Überstands, wodurch man eine rekombinante Pilz-Cellulase-Zusammensetzung erhält.
- 40 10. Verfahren gemäß Anspruch 9, wobei nach der Filtration ein antimikrobielles Mittel zu der rekombinanten Pilz-Cellulase-Zusammensetzung gegeben wird.
- 45 11. Verfahren gemäß mindestens einem der Ansprüche 8 bis 10, ferner umfassend den Schritt des Reinigens eines Expressionsprodukts aus der isolierten rekombinanten Pilz-Cellulase-Zusammensetzung.
- 50 12. Pilz-Cellulase-Zusammensetzung, abgeleitet aus einem filamentösen Pilz, umfassend eine rekombinante veränderte extrazelluläre  $\beta$ -Glucosidase, hergestellt durch das Verfahren von mindestens einem der Ansprüche 8 bis 10 nur zu dem Ausmaß, in welchem sie abhängig von Anspruch 1(b) sind.
- 55 13. Verfahren zur Herstellung von Glucose aus Cellulose-Arten oder Heteroglykanen, umfassend die Verwendung einer rekombinanten Pilz-Cellulase-Zusammensetzung mit erhöhter  $\beta$ -Glucosidase-Aktivität, hergestellt durch das Verfahren von mindestens einem der Ansprüche 8 bis 10, nur zu dem Ausmaß, in dem sie abhängig von Anspruch 1(a) sind.
14. Verfahren zum Abbau von Cellulose-Arten, vorhanden in Einsatzmaterial, Biomasse oder Bodensatz-Schlamm, umfassend die Verwendung einer rekombinanten Pilz-Cellulase-Zusammensetzung mit erhöhter  $\beta$ -Glucosidase-Aktivität, hergestellt durch das Verfahren von mindestens einem der Ansprüche 8 bis 10, nur zu dem Ausmaß, in dem sie abhängig von Anspruch 1(a) sind.
15. Verfahren zum Abbau von Cellulose zu Glucose, umfassend die Verwendung einer rekombinanten Pilz-Cellulase-Zusammensetzung mit erhöhter  $\beta$ -Glucosidase-Aktivität, hergestellt durch das Verfahren von mindestens einem



der Ansprüche 8 bis 10, nur zu dem Ausmaß, in dem sie abhängig von Anspruch 1(a) sind.

16. Verfahren zur Verwendung der rekombinanten Pilz-Cellulase-Zusammensetzung mit erhöhter  $\beta$ -Glucosidas -Aktivität, hergestellt durch das Verfahren von mindestens einem der Ansprüche 8 bis 10, nur zu dem Ausmaß, in dem sie abhängig von Anspruch 1(a) sind, oder gereinigter  $\beta$ -Glucosidase, hergestellt aus dem Verfahren nach Anspruch 12, in Nahrungsmittelzusätzen, um Geschmack und Aroma zu verstärken.
17. Reinigungsmittelzusammensetzung, umfassend einen reinigungswirksamen Gehalt eines Tensids und mindestens 0,0001 Gew.-% einer rekombinanten Pilz-Cellulase-Zusammensetzung mit erhöhter  $\beta$ -Glucosidase-Aktivität, hergestellt durch das Verfahren von mindestens einem der Ansprüche 8 bis 10, nur zu dem Ausmaß, in dem sie abhängig von Anspruch 1(a) sind.
18. Verfahren zur Verwendung der Cellulase-Zusammensetzung mit erhöhter  $\beta$ -Glucosidase-Aktivität, hergestellt durch das Verfahren von mindestens einem der Ansprüche 8 bis 10, nur zu dem Ausmaß, daß sie abhängig von Anspruch 1(a) sind, wobei das Verfahren das Zugeben der rekombinanten Pilz-Cellulase-Zusammensetzung zu Hefe und Biomasse unter Bildung einer Mischung und das Kultivieren der Mischung bei einer ausreichenden Temperatur und während einer ausreichenden Zeitdauer umfaßt, um Ethanol herzustellen.
19. Transformanten, hergestellt durch das Verfahren gemäß Anspruch 7 nur zu dem Ausmaß, zu dem es abhängig von Anspruch 1(a) ist.
20. Verfahren zur Verwendung von Transformanten, welche erhöhte extrazelluläre  $\beta$ -Glucosidase-Aktivität aufweisen, von Anspruch 19, nur zu dem Ausmaß, zu dem es abhängig von Anspruch 1(a) ist, wobei das Verfahren das Zugeben der Transformanten zu Hefe und Biomasse unter Bildung einer Mischung und das Kultivieren der Mischung bei einer ausreichenden Temperatur und während einer ausreichenden Zeitdauer umfaßt, um Ethanol herzustellen.
21. Verfahren gemäß Anspruch 18 oder Anspruch 20, wobei die Hefe B. clausenii, S. cerevisiae, Cellulolyticus acidothermophilum, C. brassicae, C. lustinariae, S. uvarum oder Schizosaccharomyces pombe ist.
22. Nukleotidsequenz eines bgl1-Gens, wobei die gesamte Sequenz oder ein Teil markiert ist oder nicht, zur Verwendung als eine Sonde, wobei das bgl1-Gen die Nukleotidsequenz von Figur 1 aufweist.
23. Verfahren zur Verwendung einer Sonde, umfassend eine Nukleotidsequenz nach Anspruch 22, zum Identifizieren eines  $\beta$ -Glucosidase-Gens aus einem filamentösen Pilz.
24. Verfahren nach Anspruch 23, wobei die Sonde aus Trichoderma reesei abgeleitet ist.
25. Verfahren nach Anspruch 23 oder Anspruch 24, wobei eine oder mehrere Sonden verwendet werden können, um das  $\beta$ -Glucosidase-Gen aus einem filamentösen Pilz zu identifizieren.

#### Revendications

1. Procédé pour modifier l'expression de la  $\beta$ -glucosidase extracellulaire dans un champignon filamenteux, comprenant la transformation dudit champignon à l'aide d'un vecteur d'expression contenant une séquence d'ADN fongique qui :
  - (a) est capable d'améliorer l'expression de la  $\beta$ -glucosidase par la présence d'au moins une copie supplémentaire d'un gène de la  $\beta$ -glucosidase fongique ; ou
  - (b) code une  $\beta$ -glucosidase extracellulaire modifiée, c'est-à-dire une enzyme possédant une séquence d'acides aminés qui a été modifiée, par rapport à celle qui est codée par le gène bgl1 dérivé de Trichoderma reesei, par manipulation de ladite séquence d'ADN de bgl1.
2. Procédé pour améliorer l'expression de la  $\beta$ -glucosidase extracellulaire selon la revendication 1(a), dans lequel ledit vecteur d'expression comprend la totalité d'une région codante d'un gène de  $\beta$ -glucosidase fongique et les séquences nécessaires pour la transcription et la traduction du gène de la  $\beta$ -glucosidase.

3. Procédé selon la revendication 1 ou la revendication 2, dans lequel ledit champignon filamenteux est sélectionné parmi les genres Trichoderma, Aspergillus, Neurospora, Humicola et Penicillium.
- 5 4. Procédé selon la revendication 3, dans lequel ledit champignon filamenteux est Trichoderma reesei, Trichoderma viridae, Trichoderma koningii, Aspergillus niger, Aspergillus nidulans, Aspergillus wentii, Aspergillus oryzae, Aspergillus phoenicis, Neurospora crassa, Humicola grisea, Penicillium pinophilum ou Penicillium oxalicum.
- 10 5. Procédé selon l'une quelconque des revendications précédentes, dans lequel ledit gène de la  $\beta$ -glucosidase est un gène bgl1 dérivé de Trichoderma reesei.
6. Procédé selon la revendication 5, dans lequel ledit gène bgl1 comprend les acides aminés codés par la séquence nucléotidique allant de 311 à 2.679 dans la Figure 1.
- 15 7. Procédé selon l'une quelconque des revendications précédentes, comprenant, en outre, l'étape qui consiste à isoler les transformants qui expriment la  $\beta$ -glucosidase modifiée.
8. Procédé selon la revendication 7, comprenant, en outre, les étapes qui consistent à :
  - 20 (a) cultiver lesdits transformants dans des conditions permettant leur croissance ; et
  - (b) isoler une composition de cellulase fongique recombinante produite par lesdits transformants.
9. Procédé selon la revendication 8, dans lequel on isole ladite composition de cellulase fongique recombinante en :
  - 25 (a) centrifugeant ledit milieu de culture contenant lesdits transformants de manière à former un surnageant et un culot ; et
  - (b) filtrant ledit surnageant de manière à obtenir une composition de cellulase fongique recombinante.
- 30 10. Procédé selon la revendication 9, dans lequel on ajoute un agent antimicrobien à ladite composition de cellulase fongique recombinante après la filtration.
11. Procédé selon l'une quelconque des revendications 8 à 10, comprenant, en outre, l'étape qui consiste à purifier un produit d'expression de ladite composition de cellulase fongique recombinante isolée.
- 35 12. Composition de cellulase fongique dérivée d'un champignon filamenteux, comprenant une  $\beta$ -glucosidase extra-cellulaire modifiée recombinante produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(b).
- 40 13. Procédé pour produire du glucose à partir de dérivés de cellulose ou d'hétéroglycans, comprenant l'utilisation d'une composition de cellulase fongique recombinante possédant une activité  $\beta$ -glucosidase accrue produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(a).
- 45 14. Procédé pour dégrader les dérivés de cellulose présents dans les produits de base, la biomasse ou les boues, comprenant l'utilisation d'une composition de cellulase fongique recombinante possédant une activité  $\beta$ -glucosidase accrue produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(a).
- 50 15. Procédé pour dégrader la cellulose en glucose, comprenant l'utilisation d'une composition de cellulase fongique recombinante possédant une activité  $\beta$ -glucosidase accrue produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(a).
- 55 16. Procédé pour utiliser la composition de cellulase fongique recombinante possédant une activité  $\beta$ -glucosidase accrue produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(a), ou une  $\beta$ -glucosidase purifiée produite à partir du procédé selon la revendication 12, dans des adjuvants alimentaires pour renforcer la saveur et les arômes.
17. Composition détergente comprenant une quantité, efficace pour le nettoyage, de tensioactif et au moins 0,0001 pour cent en poids d'une composition de cellulase fongique recombinante possédant une activité  $\beta$ -glucosidase

accrue produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(a).

- 5 18. Procédé pour utiliser la composition de cellulase possédant une activité  $\beta$ -glucosidase accrue produite par le procédé selon l'une quelconque des revendications 8 à 10 seulement dans la mesure où elles sont dépendantes de la revendication 1(a), dans lequel ledit procédé comprend les étapes consistant à ajouter ladite composition de cellulase fongique recombinante à de la levure et de la biomasse afin de former un mélange, et à cultiver ledit mélange à une température suffisante et pendant une période de temps suffisamment longue pour produire de l'éthanol.
- 10 19. Transformants produits par le procédé selon la revendication 7 seulement dans la mesure où elle est dépendante de la revendication 1(a).
- 15 20. Procédé pour utiliser les transformants possédant une activité  $\beta$ -glucosidase extracellulaire accrue selon la revendication 19 seulement dans la mesure où elle est dépendante de la revendication 1(a), ledit procédé comprenant les étapes consistant à ajouter lesdits transformants à de la levure et de la biomasse afin de former un mélange, et à cultiver ledit mélange à une température suffisante et pendant une période de temps suffisamment longue pour produire de l'éthanol.
- 20 21. Procédé selon la revendication 18 ou la revendication 20, dans lequel ladite levure est B. clausenii, S. cerevisiae, Cellulolyticus acidothermophilum, C. brassicae, C. lustinaniae, S. uvarum ou Schizosaccharomyces pombe.
- 25 22. Séquence nucléotidique d'un gène bgl1 dont la séquence en entier, ou une portion de celle-ci, est ou n'est pas marquée en vue d'être utilisée en tant que sonde, dans laquelle le gène bgl1 possède la séquence nucléotidique de la Figure 1.
23. Méthode d'utilisation d'une sonde comprenant une séquence nucléotidique selon la revendication 22 en vue d'identifier un gène de la  $\beta$ -glucosidase d'un champignon filamenteux.
- 30 24. Méthode selon la revendication 23, dans laquelle ladite sonde est dérivée de Trichoderma reesei.
25. Méthode selon la revendication 23 ou la revendication 24, dans laquelle on peut utiliser une ou plusieurs sonde(s) pour identifier ledit gène de la  $\beta$ -glucosidase d'un champignon filamenteux.

35

40

45

50

55

**FIG. 1**

TTGGCCACAGA GGGAGAGTTC GGGCTACCGC TTGGTGGAGG AAATGATGGC CCAAGGGCTC	60
AAATGGTAAA TCTGGGTGTG GGTAGGAGTG CAAAGATGGG ATTGGGCGC AATGCTGCG	120
AGCGGAGTG TTTCTGCAAC GTTATCCAGG AGATTGGGC TTGCGCAAGA GGGAGTTGAC	180
GGGAGAGTC CCAACTGGTT CCTTCAGTAA GCGACCGTG GCAGACTATA TAACTTGTGG	240
ACAAGACTCT GCTTTGTGA GTTCTTCTA CCAGTCTTGA CCAAGACAT TCTGTTGAGC	300
CCAATCAGAA ATG CGT TAC CGA ACA GCA GCT GCG CTG GCA CTT GGC ACT	349
Met Arg Tyr Arg Thr Ala Ala Ala Leu Ala Leu Ala Thr	
1 5 10	
GGG CCG TTT GCT AGG GCA GAC AGT CA GTATAGCTGG TCCATCTGG	395
Gly Pro Phe Ala Arg Ala Asp Ser His	
15 20	

**FIG. 1 (Con't.)**

GATGTGATAT GTATCTGGA GACAACATOC TGACTCTTGA ATCAAGGTAG C TCA ACA Ser Thr	452
TOG GGG GGC TGG GCT GAG GCA GTT GTA OCT OCT GCA GGG ACT OCA TGG Ser Gly Ala Ser Ala Glu Ala Val Val Pro Pro Ala Gly Thr Pro Trp 25 30 35 40	500
GGA AOC GCG TAC GAC AAG GCG AAG GGC GCA TTG GCA AAG CTC AAT CTC Gly Thr Ala Tyr Asp Lys Ala Lys Ala Ala Leu Ala Lys Leu Asn Leu 45 50 55	548
CAA GAT AAG GTC GGC ATC GTG AGC GGT GTC GGC TGG AAC GGC GGT OCT Gln Asp Lys Val Gly Ile Val Ser Gly Val Gly Trp Asn Gly Gly Pro 60 65 70	596
TGC GTT GGA AAC ACA TCT CCG GGC TOC AAG ATC AGC TAT OCA TGG CTA Cys Val Gly Asn Thr Ser Pro Ala Ser Lys Ile Ser Tyr Pro Ser Leu 75 80 85	644
TGC CTT CAA GAC GGA CCC CTC GGT GTT CGA TAC TGG ACA GGC AGC ACA Cys Leu Gln Asp Gly Pro Leu Gly Val Arg Tyr Ser Thr Gly Ser Thr 90 95 100	692
GOC TTT ACG CCG GGC GTT CAA GCG GGC TGG ACG TGG GAT GTC AAT TTG Ala Phe Thr Pro Gly Val Gln Ala Ala Ser Thr Trp Asp Val Asn Leu 105 110 115 120	740
ATC CCG GAA CGT GGA CAG TTC ATC GGT GAG GAG GTG AAG GGC TGG GCG Ile Arg Glu Arg Gly Gln Phe Ile Gly Glu Glu Val Lys Ala Ser Gly 125 130 135	788
ATT CAT GTC ATA CTT GGT OCT GTG GCT GCG CCG CTG GGA AAG ACT CCG Ile His Val Ile Leu Gly Pro Val Ala Gly Pro Leu Gly Lys Thr Pro 140 145 150	836
CAG GGC GGT CCG AAC TGG GAG GGC TTC GGT GTC GAT OCA TAT CTC ACG Gln Gly Gly Arg Asn Trp Glu Gly Phe Gly Val Asp Pro Tyr Leu Thr 155 160 165	884
GGC ATT GOC ATG GGT CAA AOC ATC AAC GGC ATC CAG TGG GTA GGC GTG Gly Ile Ala Met Gly Gln Thr Ile Asn Gly Ile Gln Ser Val Gly Val 170 175 180	932
CAG GCG ACA GCG AAG CAC TAT ATC CTC AAC GAG CAG GAG CTC AAT CGA Gln Ala Thr Ala Lys His Tyr Ile Leu Asn Glu Gln Glu Leu Asn Arg 185 190 195 200	980
GAA AOC ATT TGG AGC AAC OCA GAT GAC CGA ACT CTC CAT GAG CTG TAT Glu Thr Ile Ser Ser Asn Pro Asp Asp Arg Thr Leu His Glu Leu Tyr 205 210 215	1028

**FIG. 1 (Con't.)**

ACT TGG OCA TTT GGC GAC GGG GTT CAG GGC AAT GTC GCT TCT GTC ATG Thr Trp Pro Phe Ala Asp Ala Val Gln Ala Asn Val Ala Ser Val Met 220 225 230	1076
TGC TGG TAC AAC AAG GTC AAT ACC ACC TGG GGC TGC GAG GAT CAG TAC Cys Ser Tyr Asn Lys Val Asn Thr Thr Trp Ala Cys Glu Asp Gln Tyr 235 240 245	1124
ACG CTG CAG ACT GTG CTG AAA GAC CAG CTG GGG TTC OCA GGC TAT GTC Thr Leu Gln Thr Val Leu Lys Asp Gln Leu Gly Phe Pro Gly Tyr Val 250 255 260	1172
ATG ACG GAC TGG AAC GCA CAG CAC ACG ACT GTC CAA AGC GCG AAT TCT Met Thr Asp Trp Asn Ala Gln His Thr Thr Val Gln Ser Ala Asn Ser 265 270 275 280	1220
GGG CTT GAC ATG TCA ATG OCT GGC ACA GAC TTC AAC GGT AAC AAT CCG Gly Leu Asp Met Ser Met Pro Gly Thr Asp Phe Asn Gly Asn Asn Arg 285 290 295	1268
CTC TGG GGT OCA GCT CTC ACC AAT GCG GTA AAT AGC AAT CAG GTC CCC Leu Trp Gly Pro Ala Leu Thr Asn Ala Val Asn Ser Asn Gln Val Pro 300 305 310	1316
ACG AGC AGA GTC GAC GAT ATG GTG ACT CGT ATC CTC GGC GCA TGG TAC Thr Ser Arg Val Asp Asp Met Val Thr Arg Ile Leu Ala Ala Trp Tyr 315 320 325	1364
TTG ACA GGC CAG GAC CAG GCA GGC TAT CCG TCG TTC AAC ATC AGC AGA Leu Thr Gly Gln Asp Gln Ala Gly Tyr Pro Ser Phe Asn Ile Ser Arg 330 335 340	1412
AAT GTT CAA GGA AAC CAC AAG ACC AAT GTC AGG GCA ATT GGC AGC GAC Asn Val Gln Gly Asn His Lys Thr Asn Val Arg Ala Ile Ala Arg Asp 345 350 355 360	1460
GGC ATC GTT CTG CTC AAG AAT GAC GGC AAC ATC CTG CCG CTC AAG AAG Gly Ile Val Leu Leu Lys Asn Asp Ala Asn Ile Leu Pro Leu Lys Lys 365 370 375	1508
CCC GCT AGC ATT GGC GTC GTT GGA TCT GGC GCA ATC ATT GGT AAC CAC Pro Ala Ser Ile Ala Val Val Gly Ser Ala Ala Ile Ile Gly Asn His 380 385 390	1556
GCC AGA AAC TGG CCC TGG TGC AAC GAC AAA GGC TGC GAC GAC GGG GGC Ala Arg Asn Ser Pro Ser Cys Asn Asp Lys Gly Cys Asp Asp Gly Ala 395 400 405	1604
TTG GGC ATG GGT TGG GGT TCC GGC GGC GTC AAC TAT CCG TAC TTC GTC Leu Gly Met Gly Trp Gly Ser Gly Ala Val Asn Tyr Pro Tyr Phe Val 410 415 420	1652

**FIG. 1 (Con't.)**


GCG OOC TAC GAT GGC ATC AAT ACC AGA GCG TCT TCG CAG GGC AOC CAG Ala Pro Tyr Asp Ala Ile Asn Thr Arg Ala Ser Ser Gln Gly Thr Gln 425 430 435 440	1700
GTT ACC TTG AGC AAC ACC GAC AAC ACG TOC TCA GGC GCA TCT GCA GCA Val Thr Leu Ser Asn Thr Asp Asn Thr Ser Ser Gly Ala Ser Ala Ala 445 450 455	1748
AGA GGA AAG GAC GTC GGC ATC GTC TTC ATC AOC GGC GAC TCG GGT GAA Arg Gly Lys Asp Val Ala Ile Val Phe Ile Thr Ala Asp Ser Gly Glu 460 465 470	1796
GGC TAC ATC AOC GTG GAG GGC AAC GCG GGC GAT GGC AAC AAC CTG GAT Gly Tyr Ile Thr Val Glu Gly Asn Ala Gly Asp Arg Asn Asn Leu Asp 475 480 485	1844
CCG TCG CAC AAC GGC AAT GGC CTG GTC CAG GCG GTG GGC GGT GGC AAC Pro Trp His Asn Gly Asn Ala Leu Val Gln Ala Val Ala Gly Ala Asn 490 495 500	1892
AGC AAC GTC ATT GTT GTT GTC CAC TOC GTT GGC GGC ATC ATT CTG GAG Ser Asn Val Ile Val Val Val His Ser Val Gly Ala Ile Ile Leu Glu 505 510 515 520	1940
CAG ATT CTT GCT CTT CCG CAG GTC AAG GGC GTT GTC TGG GCG GGT CTT Gln Ile Leu Ala Leu Pro Gln Val Lys Ala Val Val Trp Ala Gly Leu 525 530 535	1988
OCT TCT CAG GAG AGC GGC AAT GCG CTC GTC GAC GTG CTG TGG GGA GAT Pro Ser Gln Glu Ser Gly Asn Ala Leu Val Asp Val Leu Trp Gly Asp 540 545 550	2036
GTC AGC OCT TCT GGC AAG CTG GTG TAC AOC ATT GCG AAG AGC OOC AAT Val Ser Pro Ser Gly Lys Leu Val Tyr Thr Ile Ala Lys Ser Pro Asn 555 560 565	2084
GAC TAT AAC ACT GGC ATC GTT TOC GGC GGC AGT GAC AGC TTC AGC GAG Asp Tyr Asn Thr Arg Ile Val Ser Gly Gly Ser Asp Ser Phe Ser Glu 570 575 580	2132
GGA CTG TTC ATC GAC TAT AAG CAC TTC GAC GAC GGC AAT ATC ACG OCG Gly Leu Phe Ile Asp Tyr Lys His Phe Asp Asp Ala Asn Ile Thr Pro 585 590 595 600	2180
CGG TAC GAG TTC GGC TAT GGA CTG T GTAAGTTTGC TAACTGAAC Arg Tyr Glu Phe Gly Tyr Gly Leu 605	2225
AATCTATTAG ACAGGTTGAC TGAACGATGA CTGTGGAATG ATAG CT TAC AOC AAG Ser Tyr Thr Lys 610	2280

**FIG. 1 (Con't.)**


TTC AAC TAC TCA GGC CTC TOC GTC TTG TOG AOC GOC AAG TCT GGT OCT Phe Asn Tyr Ser Arg Leu Ser Val Leu Ser Thr Ala Lys Ser Gly Pro 615 620 625	2328
GCG ACT GCG GGC GTT GTG GCG GGA GGC GCG AGT GAT CTG TTC CAG AAT Ala Thr Gly Ala Val Val Pro Gly Gly Pro Ser Asp Leu Phe Gln Asn 630 635 640	2376
GTC GCG ACA GTC AOC GTT GAC ATC GCA AAC TCT GGC CAA GTG ACT GGT Val Ala Thr Val Thr Val Asp Ile Ala Asn Ser Gly Gln Val Thr Gly 645 650 655 660	2424
GOC GAG GTA GOC CAG CTG TAC ATC AOC TAC OCA TCT TCA GCA OOC AGG Ala Glu Val Ala Gln Leu Tyr Ile Thr Tyr Pro Ser Ser Ala Pro Arg 665 670 675	2472
AOC OCT GCG AAG CAG CTG OGA GGC TTT GOC AAG CTG AAC CTC ACG OCT Thr Pro Pro Lys Gln Leu Arg Gly Phe Ala Lys Leu Asn Leu Thr Pro 680 685 690	2520
GGT CAG AOC GGA ACA GCA ACG TTC AAC ATC OGA OGA OGA GAT CTC ACG Gly Gln Ser Gly Thr Ala Thr Phe Asn Ile Arg Arg Arg Asp Leu Ser 695 700 705	2568
TAC TGG GAC ACG GCT TOG CAG AAA TGG GTG GTG GCG TOG GGG TOG TTT Tyr Trp Asp Thr Ala Ser Gln Lys Trp Val Val Pro Ser Gly Ser Phe 710 715 720	2616
GGC ATC ACG GTG GGA GCG ACG ACG GCG GAT ATC ACG CTG ACG ACG ACT Gly Ile Ser Val Gly Ala Ser Ser Arg Asp Ile Arg Leu Thr Ser Thr 725 730 735 740	2664
CTG TOG GTA GCG TAGGCGAGG AGGGTGAAGG GGGTTGAOCT GTGACTGTGA Leu Ser Val Ala 745	2716
GTGAGGACCG AAGGTGGGAT GCGGTGAATA CTGCAGGAAT ACAATCTTCA GGATAGGCAT	2776
CAGAGCAGTA ACATGAATGA TGAAGACGGC CGAAGCAGAA GTGAATTGAG GAGGTAGTGA	2836
TGATGAAATG TGAGGGAAGA GAGATGTTCA ATCAOCTTGT TOGAGGGAAG CTGCAAATTG	2896
GGOCTCAOCT CATCTGGCAG AGAGAAGGAA CTCTTGACAGC AGGAGTTCTG CTCACTGAGA	2956
AGAAGGCOOG GGTTAGGTC GGGCTCTTTC CGGACATOC TOOGCTOOGG CACTGTGCTG	3016
TCAAACCTGGC ACCAACA	3033







**FIG. 3A**



**FIG. 3B**






FIG. 4

A  
 $\Delta 12$   
 $\Delta 36$   
P37



6.0Kb

B  
P37  
 $\Delta 36$   
 $\Delta 12$



2.4Kb

FIG. 5A

FIG. 5B




FIG. 5C

1 2 3 4 5 7 8 9



FIG. 6




FIG. 7







FIG. 8







FIG. 10A



*Humicula grisea*

**FIG. 10B**